



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



# Increasing interdependence of multivariate distributions <sup>☆</sup>

Margaret Meyer <sup>a,\*</sup>, Bruno Strulovici <sup>b</sup>

<sup>a</sup> *Nuffield College and Dept. of Economics, Oxford University, United Kingdom*

<sup>b</sup> *Dept. of Economics, Northwestern University, USA*

Received 24 April 2010; final version received 22 July 2011; accepted 1 September 2011

Available online 10 September 2011

---

## Abstract

Orderings of interdependence are useful in many economic contexts: in assessing ex post inequality under uncertainty; in comparing multidimensional inequality; in valuing portfolios of assets or insurance policies; and in assessing systemic risk. We explore five orderings of interdependence for multivariate distributions: greater weak association, the supermodular ordering, the convex-modular ordering, the dispersion ordering, and the concordance ordering. For two dimensions, all five are equivalent, whereas for three dimensions, the first four are strictly ranked and the last two are equivalent, and for four or more dimensions, all five are strictly ranked. For the special case of binary random variables, we establish some equivalences among the orderings.

© 2011 Elsevier Inc. All rights reserved.

*JEL classification:* D63; D81; G11; G22

*Keywords:* Dependence ordering; Stochastic orders; Supermodularity; Weak association; Concordance

---



---

<sup>☆</sup> We thank Larry Epstein, Ian Jewitt, Paul Klemperer, and Jeremy Large, as well as seminar audiences at Oxford, ESSET Gerzensee, and the Paris Tech-JET Conference on Inequality and Risk. We are very grateful to the referees and the Associate Editors for their very helpful suggestions.

\* Corresponding author.

*E-mail addresses:* [margaret.meyer@nuffield.ox.ac.uk](mailto:margaret.meyer@nuffield.ox.ac.uk) (M. Meyer), [b-strulovici@northwestern.edu](mailto:b-strulovici@northwestern.edu) (B. Strulovici).

## 1. Introduction

### 1.1. Motivation

In many economic contexts, it is of interest to know whether one set of random variables displays a greater degree of interdependence than another. This paper explores several orderings of interdependence for multivariate distributions and establishes the relationships among the orderings, both in general as well as in the important special case when all random variables are binary.

Orderings of interdependence are applicable in several welfare-economic contexts. In many group settings where individual outcomes (e.g. rewards) are uncertain, members of the group may be concerned, *ex ante*, about how unequal their *ex post* rewards will be (Meyer and Mookherjee [40]; Ben-Porath et al. [9]; Gajdos and Maurin [26]; Adler and Sanchirico [2]; Chew and Sagi [17]). (This concern is distinct from concerns about the mean level of rewards and about their riskiness.) Comparisons of reward schemes then require comparisons of the degree of interdependence of the random rewards.

Another welfare-economic application concerns comparisons of inequality when individual-level data are available on multiple attributes such as income, health, and education (Atkinson and Bourguignon [6]; Bourguignon and Chakravarty [11]; Atkinson [5]). As long as the function aggregating the different attributes into an overall measure of welfare or deprivation for individuals is not additively separable across attributes, comparisons of multidimensional inequality will necessitate comparisons of the degree of interdependence among the attributes.<sup>1</sup>

Interdependence orderings are also useful for comparing the efficiency of matching mechanisms. In many matching contexts, perfectly assortative matching would be efficient and would correspond to a “perfectly positively dependent” joint distribution of the random variables representing quality in each category (dimension). When, however, matches are formed based only on noisy or coarse information (McAfee [38]), or when search is costly (Shimer and Smith [51]), or when signaling is constrained by borrowing constraints (Fernandez and Gali [25]), perfectly assortative matching will generally not arise. Fernandez and Gali [25] and Meyer and Rothschild [41] apply bivariate dependence orderings to compare the performance of different matching institutions.<sup>2</sup>

Orderings of interdependence can also be applied to compare the degree of alignment in the preferences of members of decision-making groups (Boland and Proschan [10]; Baldiga and Green [7]). Relatedly, in a strategic model of consensus building within a committee, Caillaud and Tirole [13] study how the degree of interdependence of members’ *ex ante* uncertain payoffs from a proposal affects the proposer’s optimal persuasion strategy.

In theoretical or experimental studies of information transmission in networks (Calvo-Armengol and Jackson [14], Choi, Gale, and Kariv [19]), interdependence orderings can be used

---

<sup>1</sup> There are important formal differences between these two welfare-economic applications. In the former application, the number of dimensions represents the number of individuals in the group, whereas in the latter, the number of dimensions represents the number of attributes.

<sup>2</sup> In a related application, Prat [47] explores how the composition of employee teams affects interdependence in (*ex ante* random) decisions of team members and shows how properties of the production function translate into preferences over team composition.

to assess how the degree of concordance of individuals' ex ante uncertain actions or outcomes varies with time and with social structure, as represented by the network.

Finally, in finance and insurance, valuing portfolios of assets or insurance policies requires assessing the degree of interdependence among asset returns or insurance claims (Müller and Stoyan [46]; Denuit et al. [21]). Financial economists and macroeconomists are, moreover, increasingly interested in measures and comparisons of systemic risk in financial and economic systems that reflect the interdependence in the returns of different financial institutions or sectors (Acharya [1]; Allen, Babus, and Carletti [3]; Beale et al. [8]; Hennessy and Lapan [30]).

For the special case of bivariate distributions, economists and statisticians have shown that two intuitive concepts of greater interdependence are in fact equivalent. Suppose we are comparing the degree of interdependence between  $(Y_1, Y_2)$  with that between  $(X_1, X_2)$ , where for each  $i$ ,  $Y_i$  and  $X_i$  have the same marginal distribution. The first concept is *lower-orthant dominance*. Lower-orthant dominance of  $Y$  over  $X$  requires that for all points in the support of the random vectors  $Y$  and  $X$ , the cumulative distribution function (cdf) of  $Y$  be at least as large as the cdf of  $X$ : this captures the requirement that the components of  $Y$  are more likely than those of  $X$  to both be “low” together, for any thresholds determining the precise meaning of “low”.<sup>3</sup> The second concept of greater interdependence is *supermodular dominance*, which requires that  $Ew(Y)$  be at least as large as  $Ew(X)$  for all objective functions  $w$  that are supermodular. Supermodularity (see Topkis [53] and Section 2.3) is a natural property of an objective function to capture a preference for interdependence, since its arguments are complements, not substitutes: For a supermodular function (of two or more arguments), the effect of an increase in the value of any argument is larger, the larger are the values of the other arguments. It has been shown, in the economics literature by Levy and Paroush [37], Epstein and Tanny [23], and Atkinson and Bourguignon [6] and in the statistics literature by Tchen [52], that for two-dimensional random vectors with identical marginals, lower-orthant dominance of  $Y$  over  $X$  is equivalent to  $Ew(Y) \geq Ew(X)$  for all supermodular functions  $w$ .

Economists have, nevertheless, made very little progress in the development of orderings for comparing interdependence in multivariate, as opposed to bivariate, distributions. Given the wide variety of applications for such orderings in both theoretical and empirical work, this is disappointing and may seem, at first glance, surprising. However, as we now argue, the  $n$ -dimensional case is substantially more difficult than the 2-dimensional case, for several reasons.

First, whereas positive and negative interdependence are “mirror images” of each other in two dimensions, this symmetry breaks down for more than two dimensions. In two dimensions, for any plausible concept of positive dependence, if  $Y_1$  and  $Y_2$  are positively dependent, then  $-Y_1$  and  $Y_2$  are negatively dependent. Moreover, for  $Y_1$  and  $Y_2$  with identical uniform marginals on  $[0, 1]$ , perfect positive dependence corresponds to  $Y_2 = Y_1$ , while perfect negative dependence corresponds to  $Y_2 = 1 - Y_1$ . For more than two dimensions, however, there is in general no simple way to convert a positively interdependent random vector  $(Y_1, Y_2, \dots, Y_n)$  into a negatively interdependent one. And even for  $\{Y_i\}_{i=1}^n$  with identical uniform marginals on  $[0, 1]$ , while  $Y_1 = Y_2 = \dots = Y_n$  represents perfect positive dependence, there is no obvious definition

<sup>3</sup> Given the assumption of identical marginal distributions, lower-orthant dominance is equivalent to *upper-orthant dominance*, which requires that the components of  $Y$  be more likely than those of  $X$  to both be “high” together.

of perfect negative dependence.<sup>4</sup> Since multivariate concepts of greater and lesser interdependence should be applicable to distributions displaying either positive or negative dependence, the lack of symmetry between positive and negative dependence in  $n > 2$  dimensions complicates the development of orderings.<sup>5</sup>

Second, for  $n > 2$  dimensions, there are more distinct notions of greater interdependence than there are for  $n = 2$ . Section 2 below presents five dependence orderings and shows that, whereas for  $n = 2$ , all five are equivalent, for  $n = 3$ , four are strictly ranked and only two are equivalent, and for  $n > 3$ , all five are strictly ranked. Thus, the selection of orderings of interdependence is more complicated for multivariate distributions than for bivariate ones.

Finally, even for a given ordering of greater interdependence, determining whether two multivariate distributions can be ranked according to it may be more difficult than for bivariate distributions. For  $n = 2$ , checking lower-orthant dominance is straightforward: it requires comparing the cdf's at every point in the support. For  $n > 2$ , some of the dependence orderings we study below can be implemented in an analogous fashion, by checking a given set of inequalities for each point in the support. However, for others, there exists no set of criteria that can be applied pointwise; for implementing these orderings, more sophisticated techniques or algorithms need to be developed.

## 1.2. Outline

Section 2 presents five orderings of greater interdependence for multivariate distributions. Three of these have received some attention in the statistics and economics literatures, while two of them are new.

We begin with an ordering inspired by the dependence concept of association, a concept proposed by Esary, Proschan, and Walkup [24] and widely used in statistics.<sup>6</sup> Since the definition of association cannot be meaningfully reversed to yield a concept of negative association, we turn to the concept of *weak association* (defined by Burton, Dabrowski, and Dehling [12]) and use it to define the ordering we term *greater weak association*. A random vector  $(Y_1, \dots, Y_n)$  displays greater weak association than a random vector  $(X_1, \dots, X_n)$  if  $Y$  and  $X$  have identical marginals and, for all disjoint subsets  $A, B$  of  $\{1, \dots, n\}$  and nondecreasing functions  $r, s$ ,  $Cov(r(Y_i, i \in A), s(Y_j, j \in B)) \geq Cov(r(X_i, i \in A), s(X_j, j \in B))$ .

We then present the (multivariate) supermodular ordering, which is the natural multivariate generalization of the concept of supermodular dominance discussed above for bivariate dis-

<sup>4</sup> For the set of all distribution functions with given marginals  $F_1, F_2, \dots, F_n$ , there exist upper and lower bounds for the distribution function, termed *Fréchet bounds*. The *upper Fréchet bound* is the natural candidate for the distribution exhibiting maximal positive dependence. However, while the *lower Fréchet bound* might seem like a natural candidate for the distribution displaying maximal negative dependence, the lower Fréchet bound is not in fact a proper distribution function except in very special cases (which do not include the uniform example described in the text). See Joe [36] and Müller and Stoyan [46] for more details.

<sup>5</sup> Unfortunately, this lack of symmetry between positive and negative dependence for more than two dimensions is not always recognized. See, for example, the discussion in Galeotti et al. [27, footnote 12].

<sup>6</sup> Association, though a strong concept of dependence, is strictly weaker than affiliation, which is familiar to economists from the work of Milgrom and Weber [44], who used it to formalize positive interdependence among bidders' valuations in auctions. De Castro [16] also notes that affiliation is strictly stronger than many other concepts of positive dependence. Though association has not been widely used in economics, in a pair of recent papers Calvo-Armengol and Jackson [14], respectively [15] show that the employment statuses (respectively, wages) of individuals connected by a social network are positively dependent in the sense of association.

tributions. Meyer and Mookherjee [40] and Meyer [39] contain early proposals to use it as a multivariate dependence ordering, in the context of comparisons of ex post inequality under uncertainty. In the statistical literature, the multivariate supermodular ordering was formalized by Shaked and Shanthikumar [49].

In Section 2.4, we introduce a new dependence ordering, the *convex-modular ordering*. Greater interdependence for  $Y$  than for  $X$  according to the convex-modular ordering corresponds to greater riskiness, in the sense of Rothschild and Stiglitz [48], of any aggregate  $\sum_{i=1}^n r_i(Y_i)$  compared to  $\sum_{i=1}^n r_i(X_i)$ , for any choice of  $\{r_i\}_{i=1}^n$  nondecreasing. Convex-modular objective functions are a natural way of capturing insurance companies' preferences over the degree of dependence in the claims they face.

Section 2.5 introduces another new dependence ordering, the *dispersion ordering*. The dispersion ordering is motivated by the idea that interdependence in random vectors can be compared by comparing the dispersion of the cdf's of their order statistics, with lower dispersion (assessed using the majorization ordering of vectors<sup>7</sup>) representing greater interdependence. We show that greater interdependence for  $Y$  than for  $X$  according to the dispersion ordering corresponds to greater riskiness, in the sense of Rothschild and Stiglitz [48], of the summary statistic  $\sum_{i=1}^n I_{\{Y_i > s_i\}}$  compared to  $\sum_{i=1}^n I_{\{X_i > s_i\}}$ , for any point  $(s_1, \dots, s_n)$  in the support (where  $I$  is the indicator function). A natural application of the dispersion ordering is to the assessment of systemic risk in financial systems, where systemic risk is greater the more "positively correlated" are bank failures and where the summary statistic  $\sum_{i=1}^n I_{\{Y_i > s_i\}}$  represents the number of banks that would be solvent if bank  $i$ 's failure threshold were  $s_i$ .

Finally, we present the concordance ordering, which was formalized for multivariate distributions by Joe [35] and which combines the requirement of lower-orthant dominance with that of upper-orthant dominance.<sup>8</sup> The dispersion and concordance orderings share an important advantage relative to the other three orderings, namely that they are checkable pointwise.

Section 3 presents our main results, which establish the relationships among the five orderings just described. **Theorem 1** shows that for two dimensions, all five orderings are equivalent. **Theorem 2**, on the other hand, shows that i) for three dimensions, the first four orderings presented are strictly ranked, and the dispersion and concordance orderings are equivalent, while ii) for four or more dimensions, all five orderings are strictly ranked.

Section 4 focuses on binary random variables. Binary random variables, besides being common in theoretical, experimental, and empirical applications, also help to illuminate the structure of and relationships among the dependence orderings. We study a variety of special cases with binary random variables, highlighting equivalences among the orderings in these cases. We offer a brief conclusion in Section 5.

## 2. Orderings of greater interdependence

### 2.1. Preliminaries

We consider multivariate distributions with the same number,  $n$ , of variables and identical, finite support. Focusing on finite supports simplifies notation, avoids some uninteresting technical issues, and clarifies the underlying structure of the orderings and the relationships between them.

<sup>7</sup> See Hardy, Littlewood, and Pólya [29].

<sup>8</sup> With more than two dimensions, even if  $Y$  and  $X$  have identical marginals, upper-orthant and lower-orthant dominance are no longer equivalent.

Formally, let  $L_i$  denote the finite, totally ordered set of values taken by the  $i$ th random variable, and let  $L$  denote the Cartesian product of the  $L_i$ 's.  $L_i$  is a finite subset of  $\mathbb{R}$ , and  $L$  is a finite lattice of  $\mathbb{R}^n$  with the following partial order:  $z \leq v$  if and only if  $z_i \leq v_i$  for all  $i \in N = \{1, \dots, n\}$ . If  $l_i$  denotes the cardinality of  $L_i$ , then  $L$  has  $d = \prod_{i=1}^n l_i$  elements.

As argued by Joe [36, p. 39], a natural desideratum for an ordering of interdependence is invariance to monotonic relabelings of the elements of the support for each component. Formally, we say that the ordering  $\succsim_O$  is *invariant to monotonic relabelings of coordinates* if

$$\begin{aligned} (Y_1, \dots, Y_n) \succsim_O (X_1, \dots, X_n) \\ \Rightarrow (r_1(Y_1), \dots, r_n(Y_n)) \succsim_O (r_1(X_1), \dots, r_n(X_n)) \end{aligned} \tag{1}$$

whenever  $r_i : \mathbb{R} \rightarrow \mathbb{R}$ ,  $i \in \{1, \dots, n\}$ , are all nondecreasing. This invariance property captures the idea that dependence orderings should be ordinal in that they should compare the extent to which higher (respectively, lower) realizations in one dimension are accompanied by higher (respectively, lower) realizations in other dimensions, regardless of the cardinal scales used for each dimension. An example of monotonic relabeling of coordinates is a componentwise coarsening of the support, in which some adjacent elements in the support of each component are combined. For example, starting with the lattice  $L = \{0, 1, 2\}^n$ , the nondecreasing transformations  $r_i$  with, for all  $i$ ,  $r_i(0) = 0$  and  $r_i(1) = r_i(2) = 1$  coarsen the lattice to  $\{0, 1\}^n$  by combining the realizations  $\{1, 2\}$  in each dimension.

All of the interdependence orderings we consider will be invariant to monotonic relabelings of coordinates, as defined in (1). This is straightforward to confirm from their formal definitions below. Consistent with our definition of invariance in (1), in the formal definitions of the orderings, when we consider monotonic functions, we allow these functions to be nondecreasing rather than requiring them to be increasing. None of our results is affected by this choice, but proofs are simplified by the ability to use nondecreasing indicator functions (denoted by  $I$ ).

While all of our interdependence orderings satisfy the invariance property in (1), some widely used orderings, such as the (bivariate) linear correlation coefficient, fail to satisfy this invariance and, a fortiori, are not invariant to a coarsening of the support. A higher value of the linear correlation coefficient for one bivariate empirical distribution than another is therefore not a robust indicator of greater dependence in the former distribution, since the ranking might be reversed by aggregation of the data. To see this, for  $L = \{l, m, h\}^2$ , where  $l < m < h$ , let  $(Y_1, Y_2)$  have distribution  $g$ , where  $g(l, m) = g(m, l) = g(m, m) = g(h, h) = \frac{1}{4}$ , and let  $(X_1, X_2)$  have distribution  $f$ , where  $f(l, l) = f(m, m) = f(m, h) = f(h, m) = \frac{1}{4}$ . Then  $\text{corr}(Y_1, Y_2) > (<) \text{corr}(X_1, X_2)$  if  $\frac{(l+h)}{2} > (<) m$ . This in turn implies that if the support  $L$  is coarsened by combining the realizations  $l$  and  $m$  in each dimension, then  $\text{corr}(Y_1, Y_2) > \text{corr}(X_1, X_2)$ , while if instead  $m$  and  $h$  are combined, then  $\text{corr}(Y_1, Y_2) < \text{corr}(X_1, X_2)$ .

Given that the orderings we consider below all satisfy the invariance property in (1), it is without loss of generality, and convenient notationally, to assume henceforth that the sets  $L_i$  of values taken by the  $i$ th random variable each have the form  $\{0, 1, \dots, l_i - 1\}$ .

The lattice structure of the support  $L$  and its corresponding order are useful for comparing distributions. One may label the  $d$  elements (or “nodes”) of  $L$  and view real functions on  $L$  as vectors of  $\mathbb{R}^d$ , where each coordinate of the vector corresponds to the value of the function at a specific node of  $L$ . Similarly, a multivariate distribution whose support is  $L$  can be represented as an element of the unit simplex  $\Delta_d$  of  $\mathbb{R}^d$ . For any function  $w : L \rightarrow \mathbb{R}$  and distribution  $f \in \Delta_d$ , the expected value of  $w$  given  $f$  is then the scalar product of  $w$  with  $f$ , seen as vectors of  $\mathbb{R}^d$ :

$$E[w|f] = \sum_{z \in L} w(z)f(z) = w \cdot f,$$

where  $\cdot$  denotes the scalar product of  $w$  and  $f$  in  $\mathbb{R}^d$ .

Suppose the random vectors  $Y$  and  $X$  have distributions  $g$  and  $f$ , respectively, and the former distribution dominates the latter according to some ordering “O”. We will use the phrases “ $Y$  dominates  $X$  according to the ordering O” and “ $g$  dominates  $f$  according to the ordering O” interchangeably, and we denote the former by  $Y \succ_O X$  and the latter by  $g \succ_O f$ .

Given two distributions  $g$  and  $f$  on  $L$ , define the difference between them by

$$\delta \equiv g - f. \tag{2}$$

Most (though not all) of the orderings we consider are *difference-based orderings*, in that whether or not two distributions  $g$  and  $f$  satisfy the ordering depends only on the corresponding  $\delta$ . We will frequently exploit this convenient property of difference-based orderings.<sup>9</sup>

## 2.2. Greater weak association

We begin with an ordering of greater interdependence that is inspired by the dependence concept of association. Esary, Proschan, and Walkup [24] defined association as follows:

**Definition 1 (Association).** A random vector  $Y$  with support  $L$  is **associated** if for all nondecreasing functions  $r, s : L \rightarrow \mathbb{R}$ ,  $Cov(r(Y), s(Y)) \geq 0$ .

A concept of negative association defined by reversing the inequality in the definition above would be uninteresting:  $Y$  could only be negatively associated in this strong sense if it were constant (to see this, consider functions  $r = s$ ). This is an important motivation for studying a less stringent concept, *weak association*, defined by Burton, Dabrowski, and Dehling [12], along with its negative counterpart, *negative association*, defined by Joag-Dev and Proschan [34]. In contrast to association, which allows both  $r$  and  $s$  to depend on the entire vector  $Y$ , weak association and negative association restrict them to depend on *disjoint* components of  $Y$ .<sup>10</sup>

**Definition 2 (Weak association and negative association).** A random vector  $Y$  with support  $L = \times_{i=1}^n L_i$  is **weakly associated (respectively, negatively associated)** if for any pair  $(A, B)$  of disjoint subsets of  $\{1, \dots, n\}$  and nondecreasing functions  $r : \times_{i \in A} L_i \rightarrow \mathbb{R}$  and  $s : \times_{j \in B} L_j \rightarrow \mathbb{R}$ ,  $Cov(r(Y_i, i \in A), s(Y_j, j \in B)) \geq 0$  (respectively,  $Cov(r(Y_i, i \in A), s(Y_j, j \in B)) \leq 0$ ).

We can now define an interdependence ordering corresponding to weak association and negative association as follows.<sup>11</sup>

**Definition 3 (Greater weak association).**  $Y$  displays **greater weak association** than  $X$ , denoted by  $Y \succ_{GWA} X$ , if they have identical univariate marginal distributions and for all disjoint subsets

<sup>9</sup> Some of the orderings of interdependence considered in this paper can be modified so they are responsive not only to the interdependence of the elements of a random vector but also to the levels of the elements. For clarity of focus, we will not explicitly consider these modifications of the orderings or the relations between them.

<sup>10</sup> It is clear from the definitions that association is a stronger concept than weak association. Hu, Müller, and Scarsini [31] have shown that it is strictly stronger, even for two dimensions.

<sup>11</sup> In the insurance literature, this ordering has been termed the *correlation order*—see Denuit et al. [21].



$A, B$  of  $\{1, \dots, n\}$  and nondecreasing functions  $r : \times_{i \in A} L_i \rightarrow \mathbb{R}$  and  $s : \times_{j \in B} L_j \rightarrow \mathbb{R}$ ,

$$\text{Cov}(r(Y_i, i \in A), s(Y_j, j \in B)) \geq \text{Cov}(r(X_i, i \in A), s(X_j, j \in B)).$$

Note that Definition 3 assumes that the random vectors being compared have identical univariate marginals; this is not an implication of the condition of greater covariance per se. In Section 3, we show that for  $n \geq 3$ , greater weak association is not a difference-based ordering, though Theorem 1 in Section 3 implies that for the special case of  $n = 2$ , it is.<sup>12</sup>

The greater weak association ordering has the desirable feature that  $(Y_1, \dots, Y_n)$  are weakly associated if and only if  $Y$  displays greater weak association than its *independent counterpart*, defined as the random vector  $X$  such that  $(X_1, \dots, X_n)$  are independent and, for each  $i$ ,  $X_i$  and  $Y_i$  have the same distribution.<sup>13</sup> It might seem tempting to define  $Y$  as displaying “greater association” than  $X$  if  $Y$  and  $X$  have identical marginals and for all nondecreasing functions  $r$  and  $s$  defined on  $L$ ,  $\text{Cov}(r(Y), s(Y)) \geq \text{Cov}(r(X), s(X))$ . However, such a definition would have the unappealing consequence that an associated random vector  $Y$  would not necessarily display greater association than its independent counterpart.<sup>14</sup>

### 2.3. The supermodular ordering

For any  $z, v \in L$ , denote by  $z \wedge v$  the componentwise minimum of  $z$  and  $v$ , i.e., the element of  $L$  such that for each  $i$ ,  $(z \wedge v)_i = \min\{z_i, v_i\} \in L_i$ . Let  $z \vee v$  similarly denote the componentwise maximum of  $z$  and  $v$ . A function  $w$  is said to be *supermodular* (on  $L$ ) if  $w(z \wedge v) + w(z \vee v) \geq w(z) + w(v)$  for all  $z, v \in L$ . Denote by  $e_i$  the unit vector in the  $i$ th dimension. Supermodular functions are characterized by the following property (see Topkis [53]):

$$w(z + e_i + e_j) + w(z) \geq w(z + e_i) + w(z + e_j) \tag{3}$$

for all  $i \neq j$  and  $z$  such that  $z + e_i + e_j \in L$ .<sup>15</sup>

**Definition 4** (*Supermodular ordering*). Let the random vectors  $Y$  and  $X$  have distributions  $g$  and  $f$ , respectively. The distribution  $g$  dominates the distribution  $f$  according to the **supermodular ordering**, written as  $g \succ_{SPM} f$ , if and only if  $E[w|g] \geq E[w|f]$  for all supermodular functions  $w$ .

It is clear from the definition that the supermodular ordering is a difference-based ordering.

<sup>12</sup> To verify the claim for  $n = 2$  directly, note that the only non-trivial partition of  $\{1, 2\}$  into disjoint subsets is  $\{1\}, \{2\}$ . Then given Definition 3's requirement that  $Y$  and  $X$  have identical marginals,  $\text{Cov}(r(Y_1), s(Y_2)) - \text{Cov}(r(X_1), s(X_2)) = E[r(Y_1)s(Y_2)] - E[r(X_1)s(X_2)]$ , and the sign of the right-hand side depends only on the difference between the distributions of  $Y$  and  $X$ .

<sup>13</sup> Similarly,  $(Y_1, \dots, Y_n)$  are negatively associated if and only if the independent counterpart of  $Y$  displays greater weak association than  $Y$ .

<sup>14</sup> For the support  $\{0, 1, 2\}^2$ , let  $Pr(Y_1 = Y_2 = 0) = Pr(Y_1 = Y_2 = 1) = Pr(Y_1 = Y_2 = 2) = \frac{1}{4}$  and  $Pr(Y_1 = 0, Y_2 = 2) = Pr(Y_1 = 2, Y_2 = 0) = \frac{1}{8}$ . Then it can be checked that  $Y$  is associated. However, consider the nondecreasing functions  $r(Y_1, Y_2) = I_{\{Y_1 \geq 1, Y_2 \geq 1\}}$  and  $s(Y_1, Y_2) = I_{\{Y_1 = 2 \text{ or } Y_2 = 2\}}$ . It is easy to confirm that  $\text{Cov}(r(Y_1, Y_2), s(Y_1, Y_2))$  is strictly smaller for  $Y$  than for its independent counterpart.

<sup>15</sup> For functions  $w$  defined on  $\mathbb{R}^n$  and twice differentiable, an equivalent characterization is:  $w$  is supermodular if and only if  $\frac{\partial^2 w}{\partial z_i \partial z_j} \geq 0$  for all  $z \in \mathbb{R}^n$  and all  $i \neq j$ .

To see most clearly the appeal of the supermodular ordering as an ordering of greater interdependence, consider two distributions  $g$  and  $f$  such that, for some  $z \in L$  such that  $z + e_i + e_j \in L$ , the difference  $\delta \equiv g - f$  satisfies

$$\delta(z) = \delta(z + e_i + e_j) = -\delta(z + e_i) = -\delta(z + e_j) = \alpha \tag{4}$$

for some  $\alpha > 0$ , and such that  $\delta(v) = 0$  for all other nodes  $v$  of  $L$ . In such a case, we say the distribution  $g$  is obtained from  $f$  by an *elementary transformation* (ET) of size  $\alpha$  on  $L$  which leaves unchanged the probability of all nodes other than  $z, z + e_i, z + e_j$ , and  $z + e_i + e_j$  and which raises the probability of nodes  $z$  and  $z + e_i + e_j$  by the common amount  $\alpha$ , while reducing the probability of nodes  $z + e_i$  and  $z + e_j$  by the same amount. Intuitively, such ETs increase the degree of interdependence of a multivariate distribution, as for some pair of components  $i$  and  $j$ , they make jointly high and jointly low realizations more likely, while making realizations where one component is high and the other low less likely. Furthermore, they raise interdependence without altering the marginal distribution of any component. From (3), a function  $w$  is supermodular if and only if  $w \cdot \delta \geq 0$  for any  $\delta$  of the form (4). Hence the class of supermodular functions is precisely the class for which the expectation is raised by any ET as defined in (4).

Meyer and Strulovici [42] use duality methods to characterize the supermodular ordering and develop several constructive methods to implement this characterization. In their characterization, the elementary transformations defined above play a similar role to that of mean-preserving spreads in Rothschild and Stiglitz [48] and Pigou–Dalton transfers in Atkinson [4] and Dasgupta, Sen, and Starrett [20].<sup>16</sup>

If  $g \succ_{SPM} f$ , then  $g$  and  $f$  have identical univariate marginal distributions. To see this, note that for any  $i \in \{1, \dots, n\}$  and  $k \in L_i$ , the functions  $\bar{w}(z) = I_{\{z_i \geq k\}}$  and  $\underline{w}(z) = I_{\{z_i < k\}}$  are both supermodular. Therefore  $g \succ_{SPM} f$  implies that, for each  $i \in \{1, \dots, n\}$  and  $k \in L_i$ ,

$$\begin{aligned} 0 &\leq E[\bar{w}|g] - E[\bar{w}|f] = \sum_{z: z_i \geq k} g(z) - \sum_{z: z_i \geq k} f(z) \quad \text{and} \\ 0 &\leq E[\underline{w}|g] - E[\underline{w}|f] = \sum_{z: z_i < k} g(z) - \sum_{z: z_i < k} f(z), \end{aligned} \tag{5}$$

and these inequalities together imply that  $g$  and  $f$  have identical univariate marginal distributions.

### 2.3.1. The symmetric supermodular ordering

In many contexts, it is natural to assume that the supermodular objective functions being used to compare distributions are symmetric with respect to the components of the random vectors. For example, when the function  $w$  is an ex post welfare function defined on the realized utilities of  $n$  individuals, as in the assessment of ex post inequality under uncertainty, it is natural to assume that welfare is invariant to permutations of a given  $n$ -vector of utilities over the individuals. We now formally define the symmetric supermodular ordering.

<sup>16</sup> Elementary transformations of bivariate distributions were also used by Epstein and Tanny [23] and Tchen [52] to prove the equivalence, in two dimensions, of the supermodular and lower-orthant orderings for distributions with identical marginals. Our definition of ETs in the text is more restrictive, as it requires that the four points affected by the ET be adjacent points in the lattice; this more restrictive definition allows a much simpler proof of the two-dimensional result and, more importantly, greatly facilitates the constructive methods for multivariate distributions developed in Meyer and Strulovici [42].

Call a lattice  $L = \times_{i=1}^n L_i$  symmetric if  $L_i = L_j$  for all  $i \neq j$ . Let  $\theta$  denote a real function on a symmetric lattice  $L$ . Depending on the context,  $\theta$  can represent an objective function  $w$  or a probability distribution  $f$ . We will say that the function  $\theta$  is *symmetric on  $L$*  if  $\theta(z) = \theta(\sigma(z))$  for all  $z \in L$  and for all permutations  $\sigma(z)$  of  $z$ .

**Definition 5** (*Symmetric supermodular ordering*). Let the random vectors  $Y$  and  $X$  have distributions  $g$  and  $f$ , respectively, on a symmetric lattice. The distribution  $g$  dominates the distribution  $f$  according to the **symmetric supermodular ordering**, written as  $g \succ_{SSPM} f$ , if and only if  $E[w|g] \geq E[w|f]$  for all symmetric supermodular functions  $w$ .

For an arbitrary (not necessarily symmetric) function  $\theta$ , the *symmetrized version of  $\theta$* ,  $\theta^{symm}$ , is defined as follows: for any  $z$ ,

$$\theta^{symm}(z) = \frac{1}{n!} \sum_{\sigma \in \Sigma(n)} \theta(\sigma(z)), \tag{6}$$

where  $\Sigma(n)$  is the set of all permutations of  $\{1, \dots, n\}$ . Importantly, if  $w$  is a supermodular function, then  $w^{symm}$  is supermodular. For a symmetric supermodular function  $w$ , let  $\mathcal{W}^{symm}(w)$  denote the set of supermodular functions  $\hat{w}$  on  $L$  such that the symmetrized version of  $\hat{w}$  is  $w$ , i.e.,  $\hat{w}^{symm} = w$ . Note that  $\{\mathcal{W}^{symm}(w)\}$  is a partition of the set of all supermodular functions on the symmetric lattice  $L$ . We can now state the following useful result:

**Proposition 1.** *Given a pair of distributions  $g, f$  defined on a symmetric lattice  $L$ ,  $g \succ_{SSPM} f \Leftrightarrow g^{symm} \succ_{SPM} f^{symm}$ .*

**Proof.**  $\Rightarrow$ : If for all symmetric supermodular  $w$ ,  $w \cdot g \geq w \cdot f$ , then for all symmetric supermodular  $w$ ,  $w \cdot g^{symm} \geq w \cdot f^{symm}$ . In turn, if for some symmetric supermodular  $w$ ,  $w \cdot g^{symm} \geq w \cdot f^{symm}$ , then  $\hat{w} \cdot g^{symm} \geq \hat{w} \cdot f^{symm}$  for all  $\hat{w} \in \mathcal{W}^{symm}(w)$ . Therefore, since  $\{\mathcal{W}^{symm}(w)\}_w$  partitions the set of all supermodular functions on  $L$ ,  $w \cdot g^{symm} \geq w \cdot f^{symm}$  for all symmetric supermodular  $w$  implies that  $g^{symm} \succ_{SPM} f^{symm}$ .

$\Leftarrow$ : If for all supermodular  $w$ ,  $w \cdot g^{symm} \geq w \cdot f^{symm}$ , then for all supermodular  $w$ ,  $w^{symm} \cdot g^{symm} \geq w^{symm} \cdot f^{symm}$ . This is equivalent to  $w^{symm} \cdot g^{symm} \geq w^{symm} \cdot f^{symm}$  for all symmetric supermodular  $w^{symm}$ . This in turn implies that for all symmetric supermodular  $w^{symm}$ ,  $w^{symm} \cdot g \geq w^{symm} \cdot f$ .  $\square$

**Proposition 1** states that one can characterize the symmetric supermodular ordering in terms of the supermodular order applied to symmetrized distributions.

The symmetric supermodular ordering has a very simple form for random vectors for which each component has a binary support  $\{0, 1\}$ , so the lattice is  $L = \{0, 1\}^n$ . To state the result, first define, for a random vector  $Y$  with support  $L = \{0, 1\}^n$ ,

$$c(Y) \equiv \sum_{i=1}^n I_{\{Y_i=1\}}.$$

The “count function”  $c(Y)$  gives the number of components of  $Y$  for which the realization takes the value 1. Now recall the definition of the univariate convex ordering.<sup>17</sup>

**Definition 6** (*Univariate convex ordering*). For random variables  $Z$  and  $V$  with support  $S \subseteq \mathbb{R}$ ,  $Z$  dominates  $V$  according to the **convex ordering**, written as  $Z \succ_{CX} V$ , if  $Ew(Z) \geq Ew(V)$  for all convex functions  $w : S \rightarrow \mathbb{R}$ .

Since  $w(z) = z$  and  $w(z) = -z$  are both convex functions,  $Z \succ_{CX} V$  implies  $EZ = EV$ . The convex ordering is equivalent to the ordering of greater riskiness studied by Rothschild and Stiglitz [48].

**Proposition 2.** For random vectors  $Y$  and  $X$  distributed on  $L = \{0, 1\}^n$ ,  $Y \succ_{SSPM} X$  if and only if  $c(Y) \succ_{CX} c(X)$ .

**Proof.** Any symmetric function  $w$  defined on  $L = \{0, 1\}^n$  can be written as

$$w(Y_1, \dots, Y_n) = \phi(c(Y_1, \dots, Y_n)), \tag{7}$$

for some function  $\phi : \{0, 1, \dots, n\} \rightarrow \mathbb{R}$ . Furthermore, a symmetric function  $w$  on  $\{0, 1\}^n$  is supermodular if and only if the function  $\phi(\cdot)$  in (7) is convex.  $\square$

In Sections 2.4 and 2.5, we propose two new orderings of greater interdependence.

#### 2.4. The convex-modular ordering

In many contexts, the objective functions used to compare interdependence in multivariate distributions have the form

$$w(z) = \phi(r_1(z_1) + \dots + r_n(z_n)), \quad \text{with } \phi(\cdot) \text{ convex and } \{r_i(\cdot)\}_{i=1}^n \text{ nondecreasing.}$$

We will term such functions *convex-modular*, as they take a convex transformation of a modular (i.e. additively separable) aggregate,  $\sum_{i=1}^n r_i(z_i)$ . Any convex-modular function is supermodular, and therefore the expectation of any convex-modular function is not decreased by any elementary transformation of the form defined in (4). The proof of Proposition 2 rested on the observation that any symmetric supermodular function  $w$  defined on  $L = \{0, 1\}^n$  is convex-modular, with  $\sum_{i=1}^n r_i(z_i) = \sum_{i=1}^n I_{\{z_i=1\}}$ . Convex-modular functions arise naturally in an insurance context, where  $Z$  represents a vector of losses incurred by individuals  $1, \dots, n$ , all of whom are insured by a given insurer, and where the insurance contract of individual  $i$  obliges the insurer to pay compensation  $r_i(Z_i)$ , which would take the form  $r_i(Z_i) = \min\{m_i, \max\{(1 - \beta_i)(Z_i - d_i), 0\}\}$  for a policy with a deductible  $d_i$ , a copayment rate  $\beta_i$  for the insured, and a compensation limit  $m_i$ . The total compensation paid out by the insurer is then  $\sum_{i=1}^n r_i(Z_i)$ , and the insurer is concerned with the riskiness of this total, so evaluates the cost of this payout using a convex objective function  $\phi$ .<sup>18</sup> These observations motivate us to define the following difference-based ordering:

<sup>17</sup> We do not require the support  $S$  to be convex, since we wish to apply the ordering to discrete random variables. A function defined on  $S \subset \mathbb{R}$  is convex if it can be extended to a convex function on  $\mathbb{R}$ .

<sup>18</sup> See Denuit et al. [21] for more details.

**Definition 7** (*Convex-modular ordering*). Let the random vectors  $Y$  and  $X$  have distributions  $g$  and  $f$ , respectively. The distribution  $g$  dominates the distribution  $f$  according to the **convex-modular ordering**, written as  $g \succ_{CXMOD} f$ , if and only if  $E[w|g] \geq E[w|f]$  for all convex-modular functions  $w$ .

This definition is equivalent to the requirement that  $E[w|g] \geq E[w|f]$  for all functions  $w$  that are nonnegative weighted sums of convex-modular functions. It follows from the definition that  $Y \succ_{CXMOD} X$  if and only if, for all nondecreasing  $\{r_i\}_{i=1}^n$ ,  $\sum_{i=1}^n r_i(Y_i) \succ_{CX} \sum_{i=1}^n r_i(X_i)$ .

Since  $\bar{w}(z) = I_{\{z_i \geq k\}}$  and  $\underline{w}(z) = -I_{\{z_i \geq k\}}$  are both convex-modular functions, it follows, from the same logic as for the supermodular ordering, that  $Y \succ_{CXMOD} X$  implies that  $Y$  and  $X$  have identical marginals.

It is natural to define a symmetric counterpart of the convex-modular ordering. It is clear from our definition in (6) of the symmetrized version of a function, that if  $w$  is a convex-modular function, then  $w^{symm}$  is a nonnegative weighted sum of convex-modular functions. Let  $CM^*$  denote the set of *nonnegative weighted sums of convex-modular functions* (where the dependence on a given  $L$  is implicit). It follows from the above observation that  $CM^*$  is closed under symmetrization (although the set of convex-modular functions itself is not). As a consequence, we define the symmetric convex-modular ordering as follows:

**Definition 8** (*Symmetric convex-modular ordering*). Let the random vectors  $Y$  and  $X$  have distributions  $g$  and  $f$ , respectively, on a symmetric lattice. The distribution  $g$  dominates the distribution  $f$  according to the **symmetric convex-modular ordering**, written as  $g \succ_{SCXMOD} f$ , if and only if  $E[w|g] \geq E[w|f]$  for all symmetric functions  $w \in CM^*$ .

As noted above,  $g \succ_{CXMOD} f$  if and only if  $E[w|g] \geq E[w|f]$  for all  $w \in CM^*$ . Using this equivalence, it is then straightforward to adapt the proof of Proposition 1 to show:

**Proposition 3.** *Given a pair of distributions  $g, f$  defined on a symmetric lattice  $L$ ,  $g \succ_{SCXMOD} f \Leftrightarrow g^{symm} \succ_{CXMOD} f^{symm}$ .*

### 2.5. The dispersion ordering

Another notion of greater interdependence in  $Y$  than in  $X$  reflects the idea that the distribution functions of the order statistics of  $Y$  should be “closer together” or less dispersed than the distribution functions of the order statistics of  $X$ . To understand the link between dispersion of order statistics and interdependence, suppose  $(Y_1, Y_2)$  and  $(X_1, X_2)$  both have symmetric distributions on  $\{0, 1\}^2$ , and that  $Pr(Y_i = 1) = Pr(X_i = 1) = \frac{1}{2}$ . Let  $Y_1$  and  $Y_2$  be perfectly positively dependent: the realizations  $(0, 0)$  and  $(1, 1)$  both have probability  $\frac{1}{2}$ . Let  $X_1$  and  $X_2$  be perfectly negatively dependent: the realizations  $(0, 1)$  and  $(1, 0)$  both have probability  $\frac{1}{2}$ . For the order statistics of  $Y$ ,  $\min\{Y_1, Y_2\}$  and  $\max\{Y_1, Y_2\}$ ,  $Pr(\min\{Y_1, Y_2\} = 0) = Pr(\max\{Y_1, Y_2\} = 0) = \frac{1}{2}$ , so the two order statistics have the same distribution. By contrast, for those of  $X$ ,  $Pr(\min\{X_1, X_2\} = 0) = 1$  while  $Pr(\max\{X_1, X_2\} = 0) = 0$ , so the two order statistics have distributions as different as possible in this context. The qualitative lesson of this example is that for the more dependent random vector  $Y$ , the distribution functions of the order statistics are more similar (less dispersed) than for  $X$ .

The majorization ordering of vectors can be used to formalize the notion of lower dispersion. A vector  $a$  is said to be *majorized* by a vector  $b$ , written as  $a \prec b$ , if i) the components of the

vectors have the same total sum, and ii) for all  $k$ , the sum of the  $k$  largest entries of  $a$  is weakly smaller than the sum of the  $k$  largest entries of  $b$  (see Hardy, Littlewood, and Pólya [29]). If  $a < b$ , then the components of  $a$  are less dispersed than the components of  $b$ .

Let  $Y_{(j)}$  denote the  $j$ th order statistic of  $Y$ , i.e. the  $j$ th smallest value from  $(Y_1, \dots, Y_n)$ , and define  $X_{(j)}$  similarly. Let  $F_{Y_{(j)}}$  and  $F_{X_{(j)}}$  denote the c.d.f.'s of these order statistics. For random vectors with symmetric distributions, Shaked and Tong [50] suggested the following dependence ordering.

**Definition 9** (*Symmetric dispersion ordering*). For random vectors  $Y, X$  with symmetric distributions on a symmetric lattice  $L$ , the distribution of  $Y$  dominates that of  $X$  according to the **symmetric dispersion ordering**, written as  $Y \succ_{SDISP} X$ , if the distribution functions of the order statistics of  $Y$  are less dispersed than the distribution functions of the order statistics of  $X$ , that is,

$$(F_{Y_{(1)}}(b_0), \dots, F_{Y_{(n)}}(b_0)) < (F_{X_{(1)}}(b_0), \dots, F_{X_{(n)}}(b_0)) \quad \forall b = (b_0, \dots, b_0) \in L. \quad (8)$$

We now reformulate this dependence ordering in a manner which suggests a new ordering which is both stronger and naturally applicable to asymmetric distributions. Each vector  $b = (b_0, \dots, b_0) \in L$  can be seen as generating a (componentwise) binary coarsening of the support of the random vector  $Y$  from  $L$  to  $\{0, 1\}^n$  and a corresponding coarsened version of  $Y$ ,  $Y^b$ , such that  $Y_i^b = 0$  if  $Y_i \leq b_0$  and  $Y_i^b = 1$  if  $Y_i > b_0$ . Let  $Y_{(j)}^b$  denote the  $j$ th order statistic of  $Y^b$ , i.e. the  $j$ th smallest value from  $(Y_1^b, \dots, Y_n^b) \in \{0, 1\}^n$ , and let  $F_{Y_{(j)}^b}$  denote the distribution function of this order statistic. Then a condition equivalent to (8) is

$$(F_{Y_{(1)}^b}(0), \dots, F_{Y_{(n)}^b}(0)) < (F_{X_{(1)}^b}(0), \dots, F_{X_{(n)}^b}(0)) \quad \forall b = (b_0, \dots, b_0) \in L. \quad (9)$$

For asymmetric multivariate distributions, there is no particular reason to confine attention to binary coarsenings generated by vectors with equal components  $b = (b_0, \dots, b_0)$ . We now propose a new dependence ordering, which strengthens condition (9) by requiring that it hold for every vector  $s = (s_1, s_2, \dots, s_n) \in L$ .

For a random vector  $Y$ , define its (componentwise) *binary coarsening* corresponding to the vector  $s \in L$ ,  $Y^s$ , by

$$Y^s \equiv (Y_1^s, \dots, Y_n^s) \in \{0, 1\}^n \quad \text{where } Y_i^s = \begin{cases} 0 & \text{if } Y_i \leq s_i, \\ 1 & \text{if } Y_i > s_i. \end{cases}$$

Let  $(Y_{(1)}^s, \dots, Y_{(n)}^s) \in \{0, 1\}^n$  denote the vector of order statistics of  $Y^s$ , that is,  $Y_{(j)}^s$  equals the  $j$ th smallest value from  $(Y_1^s, \dots, Y_n^s)$ . Thus,  $Y_{(j)}^s = 0$  if there are at least  $j$  values of  $i \in \{1, \dots, n\}$  such that  $Y_i \leq s_i$ . Let  $F_{Y_{(j)}^s}$  denote the distribution function of  $Y_{(j)}^s$ .

**Definition 10** (*Dispersion ordering*). For random vectors  $Y$  and  $X$  distributed on  $L$ , consider the set of all binary coarsenings of  $Y$  and  $X$ ,  $Y^s$  and  $X^s$ , respectively, corresponding to some  $s \in L$ . The distribution of  $Y$  dominates that of  $X$  according to the **dispersion ordering**, written as  $Y \succ_{DISP} X$ , if for all  $s \in L$ , the distribution functions of the order statistics of  $Y^s$  are less dispersed than the distribution functions of the order statistics of  $X^s$ , that is,

$$(F_{Y_{(1)}^s}(0), \dots, F_{Y_{(n)}^s}(0)) < (F_{X_{(1)}^s}(0), \dots, F_{X_{(n)}^s}(0)) \quad \forall s \in L. \quad (10)$$

The following proposition illuminates the appeal and convenience of the dispersion ordering<sup>19</sup> by presenting some equivalent formulations. To state it, we define, for any  $s \in L$ ,

$$c^s(Y^s) = \sum_{i=1}^n I_{\{Y_i^s=1\}} = \sum_{i=1}^n I_{\{Y_i > s_i\}}, \tag{11}$$

which counts the number of components of  $Y^s$  that equal 1, or equivalently, the number of components of  $Y$  that strictly exceed the corresponding component of  $s$ .

**Proposition 4.** *The following three conditions are equivalent:*

- i) For  $Y, X$  with support  $L$ ,  $Y \succ_{DISP} X$ .
- ii) For all  $s \in L$ ,  $Y^s \succ_{SSPM} X^s$ .
- iii) For all  $s \in L$ ,  $c^s(Y^s) \succ_{CX} c^s(X^s)$ .

**Proof.** Since for any  $s$ ,  $Y^s$  and  $X^s$  have support  $\{0, 1\}^n$ , the equivalence of ii) and iii) follows from Proposition 2 in Section 2.3.1. To show that ii) implies i), first note that since  $Y_{(j)}^s$  is the  $j$ th smallest value from  $(Y_1^s, \dots, Y_n^s)$ ,  $F_{Y_{(j)}^s}(0) = Pr(Y_{(j)}^s = 0) = E[I_{\{Y_{(j)}^s=0\}}]$  is weakly decreasing in  $j$ . Hence the majorization condition (10) can be rewritten as

$$E \left[ \sum_{j=1}^k I_{\{Y_{(j)}^s=0\}} \right] \leq E \left[ \sum_{j=1}^k I_{\{X_{(j)}^s=0\}} \right] \quad \forall k \in \{1, \dots, n\}, \forall s \in L, \tag{12}$$

with equality required for  $k = n$ . Now observe that

$$\begin{aligned} \sum_{j=1}^k I_{\{Y_{(j)}^s=0\}} &= \sum_{j=1}^k I_{\{c^s(Y^s) < n-(j-1)\}} = \sum_{j=1}^k (1 - I_{\{c^s(Y^s) \geq n-(j-1)\}}) \\ &= k - \max\{c^s(Y^s) - (n - k), 0\}. \end{aligned} \tag{13}$$

Since for any  $s \in L$ ,  $c^s(Y^s)$  is a symmetric function of  $Y^s$ , and  $\max\{z - t, 0\}$  is convex in  $z$  for any  $t \in \mathbb{R}$ ,  $\max\{c^s(Y^s) - (n - k), 0\}$  is a symmetric supermodular function of  $Y^s$  for all  $k \in \{1, \dots, n\}$ . Therefore, ii) and (13) imply that (12) holds for all  $k \in \{1, \dots, n\}$ . Setting  $k = n$  in (13) gives

$$\sum_{j=1}^n I_{\{Y_{(j)}^s=0\}} = n - c^s(Y^s), \tag{14}$$

and since both  $c^s(Y^s)$  and  $-c^s(Y^s)$  are symmetric supermodular functions of  $Y^s$ , ii) and (14) imply that for  $k = n$ , (12) holds with equality, as required. Thus ii) implies i).

<sup>19</sup> It might, at first glance, seem more natural to define the dispersion ordering by requiring (10) to hold for all  $s$  in the extended lattice  $\bar{L} \equiv \times_{i=1}^n \bar{L}_i$ , where  $\bar{L}_i \equiv \{-1, 0, 1, \dots, l_i - 1\}$ . If  $s_i = -1$ , then  $Y_i^s = 1$  with probability 1, while if  $s_i = l_i - 1$ , then  $Y_i^s = 0$  with probability 1. Thus, including coarsenings corresponding to every  $s$  in  $\bar{L}$  would make the treatment of these two types of degenerate case symmetric. Nevertheless, it is straightforward to confirm that if (10) holds for all  $s \in L$ , then it holds for all  $s \in \bar{L}$ , so the potential alternative definition is equivalent to the one used in the text.

To show that i) implies iii), first note<sup>20</sup> that for random variables  $Z$  and  $V$  with support  $\{0, 1, \dots, n\}$ ,

$$Z \succ_{CX} V \iff E(Z) = E(V) \text{ and } \forall t \in \{1, \dots, n-1\}, E[\max\{Z-t, 0\}] \geq E[\max\{V-t, 0\}]. \tag{15}$$

Given (14), the equality in (12) for  $k = n$  implies that  $E(c^s(Y^s)) = E(c^s(X^s))$ . Given (13), the weak inequality in (12) for  $k \in \{1, \dots, n-1\}$  implies that  $E[\max\{c^s(Y^s) - t, 0\}] \geq E[\max\{c^s(X^s) - t, 0\}]$  for all  $t \in \{1, \dots, n-1\}$ . Hence, for all  $s \in L$ ,  $c^s(Y^s) \succ_{CX} c^s(X^s)$ .  $\square$

**Proposition 4** provides a simple interpretation of the dispersion ordering for random vectors  $Y$  and  $X$ . Consider any coarsening of the support  $L = \times_{i=1}^n L_i$  into  $\{0, 1\}^n$  generated by, for each dimension  $i$ , classifying all values  $z_i \leq s_i$  as 0 and all values  $z_i > s_i$  as 1, for some  $s \in L$ . Then compare the distributions of the correspondingly coarsened random vectors  $Y^s$  and  $X^s$  according to the symmetric supermodular ordering, or equivalently, given **Proposition 2**, use the univariate convex ordering to compare the distributions of the random variables  $c^s(Y^s)$  and  $c^s(X^s)$  which count, respectively, the number of components of  $Y$  such that  $Y_i > s_i$  and of  $X$  such that  $X_i > s_i$ .  $Y \succ_{DISP} X$  if and only if, for any point  $s$  in the support,  $\sum_{i=1}^n I_{\{Y_i > s_i\}}$  is riskier than  $\sum_{i=1}^n I_{\{X_i > s_i\}}$  in the sense of Rothschild and Stiglitz [48].

A natural application of the dispersion ordering is to the assessment of systemic risk in financial systems, where the  $n$  dimensions represent banks and where a bank's financial health is often summarized by classifying it as either solvent or insolvent. The degree of interdependence in failures of financial institutions is of crucial importance because "as more banks fail in the same time period, the economic disruption tends to increase disproportionately" (Beale et al. [8, p. 1]). This disproportionate increase in cost, representing an aversion to positive interdependence, can be captured by a systemic cost function  $\phi(k)$ , where  $k$  is the number of banks that fail and  $\phi$  is convex. It is natural to want rankings of expected systemic cost to be robust to variations in the precise values of the failure thresholds for different banks. These considerations point towards using the dispersion ordering of the random vectors describing banks' returns to compare levels of systemic risk under different regulatory scenarios.

It is apparent from **Proposition 4** that the dispersion ordering is a difference-based ordering. Furthermore, if  $Y \succ_{DISP} X$ , then  $Y$  and  $X$  have identical univariate marginals. To see this, observe that (10) implies that for all  $s \in L$ ,

$$\sum_{j=1}^n F_{Y_{(j)}^s}(0) = \sum_{j=1}^n F_{X_{(j)}^s}(0). \tag{16}$$

Now take  $s$  to be a vector with  $i$ th component equal to  $k \in L_i$  and all other components  $j$  equal to  $l_j - 1$ . Then for any  $Y$  and  $X$  and for any  $j \neq i$ ,  $Y_j^s = X_j^s = 0$ . Hence the first  $n-1$  order statistics of both  $Y^s$  and  $X^s$  are 0, so it follows from (16) that  $F_{Y_{(n)}^s}(0) = F_{X_{(n)}^s}(0)$ . This in turn is equivalent to  $Pr(Y_i \leq k) = Pr(X_i \leq k)$ , and since this holds for all  $i$  and all  $k \in L_i$ ,  $Y$  and  $X$  have identical marginals.

It follows from **Proposition 4** that there is an easily described algorithm for checking whether or not any given pair of  $n$ -dimensional random vectors  $Y$  and  $X$  satisfy  $Y \succ_{DISP} X$ : For each  $s \in L$ , check whether or not  $c^s(Y^s) \succ_{CX} c^s(X^s)$ . Since the support of  $c^s(Y^s)$  and  $c^s(X^s)$  is

<sup>20</sup> See, for example, Jewitt [33] and Hardy, Littlewood, and Pólya [28].



$\{0, 1, \dots, n\}$ , checking  $c^s(Y^s) \succ_{CX} c^s(X^s)$  is equivalent, given (15), to checking  $n - 1$  inequalities and 1 equality. Therefore, because an equality constraint can be rewritten as two inequality constraints, to check  $Y \succ_{DISP} X$ , it is sufficient to check  $n + 1$  inequalities for each  $s \in L$ .<sup>21</sup> The fact that  $Y \succ_{DISP} X$  is checkable pointwise is an important advantage of the dispersion ordering, relative to the greater weak association, supermodular, and convex-modular orderings.

### 2.6. The concordance ordering

Another intuitively appealing notion of greater interdependence, the concordance ordering, has been formalized for multivariate distributions by Joe [35].

**Definition 11** (*Concordance ordering*). Let the random vectors  $Y$  and  $X$  have distributions  $g$  and  $f$ , respectively. The distribution  $g$  dominates the distribution  $f$  according to the **concordance ordering**, written as  $g \succ_{CONC} f$ , if and only if

$$Pr(Y \geq z) \geq Pr(X \geq z) \quad \text{and} \quad Pr(Y \leq z) \geq Pr(X \leq z) \quad \forall z \in L.$$

For any node in the support, the concordance ordering requires that the components of  $Y$ , relative to those of  $X$ , be both more likely to be all higher than at that node and more likely to be all lower than at that node.<sup>22</sup>

It is easy to see that the concordance ordering is a difference-based ordering. As is well known,  $Y \succ_{CONC} X$  implies that  $Y$  and  $X$  have identical univariate marginals.<sup>23</sup> Note that the concordance ordering shares with our dispersion ordering the desirable feature of being checkable pointwise.

## 3. Relationships among the orderings

We are now in a position to present our main results, which establish the relationships among the orderings of interdependence defined in Section 2.

**Theorem 1** (*Orderings: two dimensions*). For two dimensions, the following orderings are equivalent: greater weak association, supermodular ordering, convex-modular ordering, dispersion ordering, and concordance ordering.

**Proof.** The equivalence for  $n = 2$  between greater weak association, the supermodular ordering, and the concordance ordering is well known (see Meyer [39, Proposition 2] and Müller and Stoyan [46, Theorem 3.8.2] for references). We now prove that for  $n = 2$  the convex-modular

<sup>21</sup> In practice, many of the inequalities generated as  $s$  varies will hold trivially or be redundant. For example, for  $L = \{0, 1\}^2$ , the set of 12 inequalities that in principle need to be checked in fact reduces to only 5, corresponding to the requirements that i)  $(Y_1, Y_2)$  and  $(X_1, X_2)$  have identical marginals (4 inequalities) and ii)  $Pr(Y_1 = 0, Y_2 = 0) \geq Pr(X_1 = 0, X_2 = 0)$ .

<sup>22</sup> See Decancq [22] for a discussion of two classes of multivariate rearrangements which together underlie the concordance order.

<sup>23</sup> To see this, first take  $z \in L$  to be a vector with  $i$ th component equal to  $k$  and all other components 0. Then  $Pr(Y \geq z) \geq Pr(X \geq z)$  becomes  $Pr(Y_i \geq k) \geq Pr(X_i \geq k)$ . Similarly, taking  $z$  to be a vector with  $i$ th component equal to  $k - 1$  and all other components equal to  $l_i - 1$ ,  $Pr(Y \leq z) \geq Pr(X \leq z)$  becomes  $Pr(Y_i \leq k - 1) \geq Pr(X_i \leq k - 1)$ . Hence, for all  $i$  and  $k \in L_i$ ,  $Pr(Y_i \geq k) = Pr(X_i \geq k)$ , so  $Y$  and  $X$  have identical marginals.

and dispersion orderings are equivalent to the other three orderings by showing that  $Y \succ_{SPM} X \Rightarrow Y \succ_{CXMOD} X \Rightarrow Y \succ_{DISP} X \Rightarrow Y \succ_{CONC} X$ .

Since every convex-modular function is supermodular, the supermodular ordering implies the convex-modular ordering. The fact that the convex-modular ordering implies the dispersion ordering follows from the proof of Proposition 4 and the facts that the functions  $\max\{c^s(Y) - (2 - k), 0\}$ , for  $k \in \{1, 2\}$ , and  $-c^s(Y)$  are convex-modular. Finally, observe that by Proposition 4, if  $(Y_1, Y_2) \succ_{DISP} (X_1, X_2)$ , then for all  $s \in L$ ,  $Pr(c^s(Y^s) = 0) \geq Pr(c^s(X^s) = 0)$  and  $Pr(c^s(Y^s) = 2) \geq Pr(c^s(X^s) = 2)$ . The first of these inequalities is equivalent to  $Pr(Y_1 \leq s_1, Y_2 \leq s_2) \geq Pr(X_1 \leq s_1, X_2 \leq s_2)$ , while the second is equivalent to  $Pr(Y_1 > s_1, Y_2 > s_2) \geq Pr(X_1 > s_1, X_2 > s_2)$ . Hence  $(Y_1, Y_2) \succ_{DISP} (X_1, X_2)$  implies  $(Y_1, Y_2) \succ_{CONC} (X_1, X_2)$ .  $\square$

**Theorem 2** (Orderings: three or more dimensions).

- a) For  $n \geq 3$ , greater weak association is strictly stronger than the supermodular ordering, which is strictly stronger than the convex-modular ordering, which is strictly stronger than the dispersion ordering, which is at least as strong as the concordance ordering.
- b) For  $n = 3$ , the dispersion ordering is equivalent to the concordance ordering, whereas for  $n > 3$ , the dispersion ordering is strictly stronger than the concordance ordering.

**Proof.** i) The proof that  $Y \succ_{GWA} X$  implies  $Y \succ_{SPM} X$  is in Appendix A. Example 1 proves that this implication is strict:

**Example 1.** Let  $L = \{0, 1\}^3$  and let  $X$  and  $Y$  have distributions  $f$  and  $g$ , respectively. Let  $f(x) = \frac{1}{3}$  if  $\sum_{i=1}^3 x_i = 1$  and  $f(x) = 0$  otherwise. Let  $g(y)$  take the values  $\frac{17}{66}, \frac{11}{66}, \frac{5}{66}, \frac{1}{66}$  when  $\sum_{i=1}^3 y_i$  takes the values 0, 1, 2, 3, respectively. For these symmetric distributions, Propositions 1 and 2 together imply that  $g \succ_{SPM} f$  on  $L = \{0, 1\}^3$  if and only if  $c(Y) \succ_{CX} c(X)$  on  $\{0, 1, 2, 3\}$ . From (15),  $c(Y) \succ_{CX} c(X)$  on  $\{0, 1, 2, 3\}$  if and only if 1)  $Ec(Y) = Ec(X)$ ; 2)  $E[\max\{c(Y) - 1, 0\}] \geq E[\max\{c(X) - 1, 0\}]$ ; and 3)  $E[\max\{c(Y) - 2, 0\}] \geq E[\max\{c(X) - 2, 0\}]$ . It is easily checked that  $Ec(Y) = Ec(X) = 1$ . Condition 2) holds since  $\frac{17}{66} = 2Pr(c(Y) = 3) + Pr(c(Y) = 2) \geq 2Pr(c(X) = 3) + Pr(c(X) = 2) = 0$ , and condition 3) holds since  $\frac{1}{66} = Pr(c(Y) = 3) \geq Pr(c(X) = 3) = 0$ . Hence,  $c(Y) \succ_{CX} c(X)$  and  $g \succ_{SPM} f$ . However, for  $r(z_1) = I_{\{z_1=1\}}$  and  $s(z_2, z_3) = I_{\{z_2=z_3=1\}}$ ,  $Cov(r(Y_1), s(Y_2, Y_3)) = \frac{1}{66} - \frac{1}{3} \cdot \frac{1}{11} < 0$ , while  $Cov(r(X_1), s(X_2, X_3)) = 0$ , so  $g \not\succ_{GWA} f$  does not hold.

ii) Since every convex-modular function is supermodular,  $Y \succ_{SPM} X$  implies  $Y \succ_{CXMOD} X$ . We show that the implication is strict by providing, in Appendix B, an example of a supermodular function on  $L = \{0, 1, 2\}^3$  which cannot be written as a nonnegative weighted sum of convex-modular functions.

iii) That  $Y \succ_{CXMOD} X$  implies  $Y \succ_{DISP} X$  follows from the proof of Proposition 4 and the facts that the functions  $\max\{c^s(Y) - (n - k), 0\}$ , for  $k \in \{1, \dots, n\}$ , and  $-c^s(Y)$  are all convex-modular. Example 2 proves that this implication is strict:

**Example 2.** Let  $L = \{0, 1, 2\} \times \{0, 1\} \times \{0, 1\}$ . Since both the convex-modular and the dispersion ordering are difference-based orderings, it is sufficient to specify  $\delta \equiv g - f$ , the difference between the distributions of  $Y$  and  $X$ . Let  $\delta(z_1, z_2, z_3) = \epsilon > 0$  if  $\sum_{i=1}^3 z_i$  is even and  $\delta(z_1, z_2, z_3) = -\epsilon < 0$  if  $\sum_{i=1}^3 z_i$  is odd. It is easily checked that  $Y \succ_{CONC} X$ . As proved in

part iv) of the proof below, for  $n = 3$ ,  $Y \succ_{CONC} X \Leftrightarrow Y \succ_{DISP} X$ . However, for the convex-modular function  $w(z) = \max\{(\sum_{i=1}^3 z_i) - 2, 0\}$ , we have  $w \cdot (g - f) = -\epsilon < 0$ , so  $Y \succ_{CXMOD} X$  does not hold.

iv) To show that  $Y \succ_{DISP} X$  implies  $Y \succ_{CONC} X$ , observe that, for  $k = 1$ , the left-hand side of (12) can be rewritten as  $E[I_{\{Y_{(1)}^s=0\}}] = 1 - Pr(Y_i > s_i \forall i)$ . Therefore, (12) implies that for all  $s \in L$ ,  $Pr(Y > s) \geq Pr(X > s)$ . Similarly, it follows from the equality in (12) for  $k = n$  and the inequality for  $k = n - 1$  that, for all  $s \in L$ ,  $E[I_{\{Y_{(n)}^s=0\}}] \geq E[I_{\{X_{(n)}^s=0\}}]$ , which implies that for all  $s \in L$ ,  $Pr(Y \leq s) \geq Pr(X \leq s)$ . Hence  $Y \succ_{CONC} X$ .

Now we show that for  $n = 3$ ,  $Y \succ_{CONC} X$  implies  $Y \succ_{DISP} X$ . First, for any  $s \in L$ , showing that (12) holds with equality for  $k = n = 3$  is equivalent, given (13) and (14), to showing that

$$E\left[\sum_{i=1}^3 I_{\{Y_i \leq s_i\}}\right] = E\left[\sum_{i=1}^3 I_{\{X_i \leq s_i\}}\right],$$

and this is true, since  $Y \succ_{CONC} X$  implies that  $Y$  and  $X$  have identical marginal distributions. Given the equality in (12) for  $k = 3$ , it remains to show that for all  $s \in L$ ,

$$E[I_{\{Y_{(1)}^s=0\}}] \leq E[I_{\{X_{(1)}^s=0\}}] \quad \text{and} \quad E[I_{\{Y_{(3)}^s=0\}}] \geq E[I_{\{X_{(3)}^s=0\}}].$$

The first of these inequalities is equivalent to  $Pr(Y > s) \geq Pr(X > s)$  and the second to  $Pr(Y \leq s) \geq Pr(X \leq s)$ , as shown above in the proof that  $Y \succ_{DISP} X$  implies  $Y \succ_{CONC} X$ . Therefore, both inequalities follow from  $Y \succ_{CONC} X$ .

Example 3 proves that for  $n > 3$ ,  $Y \succ_{CONC} X$  does not imply  $Y \succ_{DISP} X$ :

**Example 3.** Let  $L = \{0, 1\}^4$ , let  $g, f$  represent the distributions of  $Y, X$ , and let  $\delta(z_1, z_2, z_3, z_4) \equiv g - f = \epsilon > 0$  if  $\sum_{i=1}^4 z_i$  is even and  $\delta(z_1, z_2, z_3, z_4) = -\epsilon < 0$  if  $\sum_{i=1}^4 z_i$  is odd. Again, it is easily checked that  $Y \succ_{CONC} X$ . For  $s = (0, 0, 0, 0)$ , consider the convex function of  $c^s(z)$  defined by  $w = \max\{c^s(z) - 2, 0\}$ . We have  $w \cdot (g - f) = -2\epsilon < 0$ , so by Proposition 4,  $Y \succ_{DISP} X$  does not hold.  $\square$

**Remark 1.** Given the close relationship shown by Proposition 4 between the dispersion ordering and the symmetric supermodular ordering, and given that both are strictly weaker than the supermodular ordering, it is natural to ask how these two orderings relate to each other. For arbitrary supports, neither implies the other; however, as we note in Section 4, in the special case of binary random variables, the dispersion ordering is strictly stronger than the symmetric supermodular ordering.

To confirm that for arbitrary supports,  $Y \succ_{DISP} X$  does not imply  $Y \succ_{SSPM} X$ , consider Example 2, used in part iii) of the proof of Theorem 2. Expand the support from  $L = \{0, 1, 2\} \times \{0, 1\} \times \{0, 1\}$  to the symmetric  $L' = \{0, 1, 2\}^3$ . For  $z \in L' \setminus L$ , let  $\delta(z) \equiv g - f = 0$  and for  $z \in L$ , define  $\delta(z)$  as in Example 2. Then  $Y \succ_{DISP} X$  but, for the symmetric supermodular function  $w$  defined in Example 2,  $w \cdot \delta = w \cdot (g - f) = -\epsilon < 0$ , so  $Y \succ_{SSPM} X$  does not hold.

To confirm that  $Y \succ_{SSPM} X$  does not imply  $Y \succ_{DISP} X$ , consider  $Y, X$  with distributions  $g, f$ , respectively, on  $L = \{0, 1\}^3$  such that  $\delta = g - f$  is given by  $\delta(0, 0, 0) = 0$ ,  $\delta(z_1, z_2, z_3) = \epsilon > 0$  if  $\sum_{i=1}^3 z_i = 1$ ,  $\delta(1, 1, 0) = -4\epsilon$ ,  $\delta(0, 1, 1) = \delta(1, 0, 1) = -\epsilon$ , and  $\delta(1, 1, 1) = 3\epsilon$ . It is easy to check that for  $k = 0, 1, 2, 3$ ,  $Pr(c(Y) = k) - Pr(c(X) = k) = 0, 3\epsilon, -6\epsilon, 3\epsilon$ , respectively, so the distribution of  $c(Y)$  is derived from the distribution of  $c(X)$  by a mean-preserving spread. Therefore,  $c(Y) \succ_{CX} c(X)$  and hence, by Proposition 2,  $Y \succ_{SSPM} X$ . However,  $P(Y_1 = 1) -$

$P(X_1 = 1) < 0$ . Hence,  $Y$  and  $X$  do not have identical marginals, and therefore  $Y \succ_{DISP} X$  does not hold.

**Remark 2.** For  $n \geq 3$ , the strongest of our interdependence orderings, greater weak association, is not a difference-based ordering. To show this, we extend [Example 1](#), used in part i) of the proof of [Theorem 2](#). For  $L = \{0, 1\}^3$ , define  $f$  and  $g$  as in [Example 1](#). Now define  $\tilde{f}(x) = \frac{1}{2}$  if  $\sum_{i=1}^3 x_i = 0$ ,  $\tilde{f}(x) = \frac{1}{6}$  if  $\sum_{i=1}^3 x_i = 1$ , and  $\tilde{f}(x) = 0$  otherwise, and let  $\tilde{g}(y)$  take the values  $\frac{50}{66}, 0, \frac{5}{66}, \frac{1}{66}$  when  $\sum_{i=1}^3 y_i$  takes the values 0, 1, 2, 3, respectively. By construction,  $\tilde{g} - \tilde{f} = g - f$ , and we showed in part i) of the proof of [Theorem 2](#) that  $g \not\succeq_{GWA} f$  does not hold. Yet (see [Appendix C](#))  $\tilde{g} \succeq_{GWA} \tilde{f}$ . Hence for  $n \geq 3$ , greater weak association is not difference-based. Since the supermodular ordering is difference-based and the distributions in this example are symmetric, it follows that even for *symmetric* distributions, greater weak association is strictly stronger than the supermodular ordering whenever  $n \geq 3$ .

#### 4. Binary random variables

In many economic contexts, the random variables whose interdependence is to be assessed are binary. Theoretical models often focus on binary action spaces or binary outcome spaces for tractability. For example, Calvo-Armengol and Jackson's [\[14\]](#) study of the effects of social networks on interdependence in individuals' employment outcomes suppressed wage variation among employed workers and focused only on whether workers were employed or unemployed. Experimental studies often focus on binary choice spaces to simplify the subjects' decision problems as well as to simplify the data analysis. For example, Choi, Gale, and Kariv [\[19\]](#), in their experimental study of the effect of network structure on social learning and the resulting interdependence among agents' decisions, focused on a decision environment with only two states of the world, two signals, and two possible actions. In empirical work, for example on multidimensional inequality, binary classifications, such as whether or not income is below the poverty line or whether or not an individual is literate, are often inevitable features of the data.

Binary random variables, besides being common, also help to illuminate the structure of and relationships among the interdependence orderings. This section studies a variety of special cases with binary random variables. Our aims are to highlight both i) equivalences among the orderings that arise in these special cases and ii) easily checkable and easily interpretable necessary and sufficient conditions for the orderings to hold.

##### 4.1. Symmetric distributions or symmetric objective functions

The following result is valid for any number  $n$  of dimensions.

#### Proposition 5.

- a) For random vectors  $Y$  and  $X$  with symmetric distributions on  $L = \{0, 1\}^n$ , the following conditions are equivalent:
- i)  $Y \succeq_{SPM} X$ ;
  - ii)  $Y \succeq_{CXMOD} X$ ;
  - iii)  $Y \succeq_{DISP} X$ ;
  - iv)  $Y \succeq_{SDISP} X$ ;
  - v)  $c(Y) \succeq_{CX} c(X)$ .

b) For random vectors  $Y$  and  $X$  distributed on  $L = \{0, 1\}^n$ , the following conditions are equivalent:

- i)  $Y \succ_{SSPM} X$ ;
- ii)  $Y \succ_{SCXMOD} X$ ;
- iii)  $c(Y) \succ_{CX} c(X)$ .

**Proof.** a) It is clear from Definitions 9 and 10 that  $Y \succ_{DISP} X$  implies  $Y \succ_{SDISP} X$ . It follows from Definition 9 and the proof of Proposition 4 that for symmetric distributions on  $L = \{0, 1\}^n$ ,  $Y \succ_{SDISP} X$  if and only if  $c(Y) \succ_{CX} c(X)$ —the only non-trivial choice of  $b = (b_0, \dots, b_0)$  in Definition 9 is  $(0, \dots, 0)$ , in which case (8) is equivalent to  $c(Y) \succ_{CX} c(X)$ . But Propositions 1 and 2 together imply that for symmetric distributions on  $L = \{0, 1\}^n$ ,  $Y \succ_{SPM} X$  if and only if  $c(Y) \succ_{CX} c(X)$ . The remaining equivalences then follow from Theorems 1 and 2.

b) Proposition 2 shows the equivalence of  $Y \succ_{SSPM} X$  and  $c(Y) \succ_{CX} c(X)$  on  $L = \{0, 1\}^n$ . Since every symmetric  $w \in CM^*$  is supermodular, i) implies ii). Since  $c(Y)$  is symmetric and convex-modular, ii) implies iii).  $\square$

For random vectors on  $L = \{0, 1\}^n$  with asymmetric distributions, the dispersion ordering is strictly stronger than the three orderings shown to be equivalent in part b) of Proposition 5. With  $s = (0, 0, \dots, 0)$ , the majorization condition (10) defining  $Y \succ_{DISP} X$  is equivalent to  $c(Y) \succ_{CX} c(X)$ , but other choices of  $s$  generate additional conditions. In particular, as shown in Section 2.5,  $Y \succ_{DISP} X$  implies that  $Y$  and  $X$  have identical marginals, whereas this is not an implication of the three orderings in part b) of Proposition 5.<sup>24</sup>

#### 4.2. Three dimensions

**Proposition 6.** For random vectors  $Y$  and  $X$  distributed on  $L = \{0, 1\}^3$ , the following conditions are equivalent:

- i)  $Y \succ_{SPM} X$ ;
- ii)  $Y \succ_{CXMOD} X$ ;
- iii)  $Y \succ_{DISP} X$ ;
- iv)  $Y \succ_{CONC} X$ ;
- v)  $c(Y) \succ_{CX} c(X)$ ;  $Y$  and  $X$  have identical marginals; and for all  $i \neq j$ ,  $Pr(Y_i = Y_j) \geq Pr(X_i = X_j)$ .

**Proof.** All of the orderings in the proposition are difference-based orderings, so it is sufficient to work with  $\delta \equiv g - f$ . For  $L = \{0, 1\}^3$ , the values of  $\delta$  at each of the 8 nodes are displayed in Fig. 1(a). Each of conditions i)–v) implies that  $Y$  and  $X$  have identical marginals (as shown in Section 2). This in turn implies that once the 4 values of  $\delta(1, 1, 1) \equiv a$ ,  $\delta(0, 1, 1) \equiv b_1$ ,  $\delta(1, 0, 1) \equiv b_2$ , and  $\delta(1, 1, 0) \equiv b_3$  are specified (corresponding to the 4 nodes marked with black dots in Fig. 1(a)), the remaining 4 values are determined, as specified in Fig. 1(a).

We first prove iv) and v) are equivalent. Given identical marginals,  $Y \succ_{CONC} X$  if and only if

$$a \geq 0, \quad a + b_k \geq 0 \quad \forall k \in \{1, 2, 3\}, \quad \text{and} \quad 2a + \sum_{i=1}^3 b_i \geq 0. \tag{17}$$

<sup>24</sup> The second example in Remark 1 following Theorem 2 is one for which  $L = \{0, 1\}^3$  and  $Y \succ_{SSPM} X$ , but  $Y$  and  $X$  do not have identical marginals, so  $Y \succ_{DISP} X$  does not hold.




Fig. 1. (a):  $\delta = g - f$  for  $L = \{0, 1\}^3$ . (b): Labels for the nodes of  $L = \{0, 1\}^3$ .

From (15),  $c(Y) \succ_{CX} c(X)$  on  $\{0, 1, 2, 3\}$  if and only if 1)  $Ec(Y) = Ec(X)$ ; 2)  $E[\max\{c(Y) - 2, 0\}] \geq E[\max\{c(X) - 2, 0\}]$ ; and 3)  $E[\max\{c(Y) - 1, 0\}] \geq E[\max\{c(X) - 1, 0\}]$ . Given identical marginals, condition 1) holds, since (using Fig. 1(a))  $Ec(Y) - Ec(X) = 3a + 2 \sum_{i=1}^3 b_i + (-3a - 2 \sum_{i=1}^3 b_i) = 0$ . Conditions 2) and 3) are equivalent, respectively, to the first and third inequalities in (17). Finally, the condition  $Pr(Y_i = Y_j) \geq Pr(X_i = X_j)$  is equivalent, given identical marginals, to  $Pr(Y_i = Y_j = 1) \geq Pr(X_i = X_j = 1)$ , and this in turn is equivalent to  $a + b_k \geq 0, k \neq i, j$ . Hence, conditions iv) and v) are equivalent.

We now provide a simple constructive proof that for  $L = \{0, 1\}^3$ ,  $Y \succ_{CONC} X$  implies  $Y \succ_{SPM} X$ . Our constructive proof<sup>25</sup> decomposes  $\delta$  into 6 elementary transformations (ETs) of the form defined in (4): for each of the 6 faces of the cube  $L = \{0, 1\}^3$ , there is one ET involving the 4 nodes on that face. Using the labels in Fig. 1(b), let the ET involving nodes  $A, B_i, B_j$ , and  $C_k$  have size  $\bar{\alpha}_{ij}$  and the ET involving nodes  $B_k, C_i, C_j$ , and  $D$  have size  $\underline{\alpha}_{ij}$ . It is easily checked that these 6 ETs sum to  $\delta$  if and only if

$$a = \bar{\alpha}_{12} + \bar{\alpha}_{13} + \bar{\alpha}_{23} \quad \text{and} \quad \forall i, j, k \in \{1, 2, 3\}, i \neq j \neq k, \quad a + b_k = \bar{\alpha}_{ij} + \underline{\alpha}_{ij}. \quad (18)$$

Now set

$$\bar{\alpha}_{ij} = a \left( \frac{a + b_k}{3a + \sum_{i=1}^3 b_i} \right) \quad \text{and} \quad \underline{\alpha}_{ij} = \left( 2a + \sum_{i=1}^3 b_i \right) \left( \frac{a + b_k}{3a + \sum_{i=1}^3 b_i} \right). \quad (19)$$

It is apparent by inspection that Eqs. (18) are satisfied and that, if the inequalities (17) defining the concordance ordering hold, then for all  $i \neq j, \bar{\alpha}_{ij} \geq 0$  and  $\underline{\alpha}_{ij} \geq 0$ . Thus,  $g \succ_{CONC} f$  implies the existence of a sequence of nonnegative ETs that sum to  $g - f$ . Since each ET raises the expectation of any supermodular function,  $g \succ_{SPM} f$ . By Theorem 2, it then follows that conditions i)–v) are all equivalent.  $\square$

<sup>25</sup> Hu, Xie, and Ruan [32, pp. 188–189] proved this in a very indirect manner using the tool of “majorization with respect to weighted trees”.

### 4.3. Four dimensions and “top-to-bottom” symmetry

The equivalence demonstrated in Proposition 6 for the three-dimensional cube between the supermodular ordering and the concordance ordering breaks down if we increase either the number of dimensions, as shown by Example 3 (used in part iv) of the proof of Theorem 2), or the number of points in the support for some dimension, as shown by Example 2 (used in part iii) of the same proof). Nevertheless, some interesting equivalences do persist in higher dimensions and can be demonstrated using a similar constructive method of proof.

Consider four-dimensional random vectors with support  $L = \{0, 1\}^4$ , and assume now that their distributions satisfy a symmetry condition we term “top-to-bottom symmetry”. We say that the distribution of a random vector  $Z$  satisfies *top-to-bottom symmetry* if for any  $a \in \{0, 1\}^4$ ,  $P(Z = a) = P(Z = (1, 1, 1, 1) - a)$ . Top-to-bottom symmetry arises naturally in a variety of settings. We give two examples: matching with frictions and social learning in networks.

In a matching context, suppose the four dimensions represent managers, supervisors, workers, and firms, and suppose that for each dimension, there is one representative (individual or firm) with high quality ( $z_i = 1$ ) and one with low quality ( $z_i = 0$ ). Production requires forming a “team” consisting of exactly one manager, one supervisor, one worker, and one firm, and the output of such a team is a supermodular function of the qualities of each of its four components. Supermodularity of the production function  $w$  implies that it would be output-maximizing for the four high-quality individuals/firm to be matched and for the four low-quality individuals/firm to be matched, in which case output would be  $w(1, 1, 1, 1) + w(0, 0, 0, 0)$ . However, informational frictions may prevent such an outcome being reached and cause the matching process to be stochastic. Nevertheless, as long as the stochastic process is certain to divide the individuals into two teams, each consisting of one representative from each dimension, the distribution over team quality vectors necessarily satisfies top-to-bottom symmetry. This is so since, if one of the teams formed has individuals with qualities  $(a_1, a_2, a_3, a_4) \in \{0, 1\}^4$ , the other team must consist of individuals with qualities  $(1 - a_1, 1 - a_2, 1 - a_3, 1 - a_4)$ , given that for each dimension, there is one representative with quality 0 and one with quality 1.

Choi, Gale, and Kariv [19] experimentally investigate the effect of network structure on the degree of interdependence of choices in social learning situations. Each agent  $i$  can choose between two actions, 0 and 1, and the randomness in the environment generates, for each period, an ex ante distribution over action vectors  $(a_1, \dots, a_n) \in \{0, 1\}^n$ . To simplify the computation of equilibrium behavior, they assume that the environment is symmetric with respect to the two states of the world, the two signals, and the two actions. Given this symmetry, in each period and for any network, the ex ante probabilities of the action vectors  $(a_1, \dots, a_n)$  and  $(1 - a_1, \dots, 1 - a_n)$  are equal, so the distributions satisfy top-to-bottom symmetry.

**Proposition 7.** *For random vectors  $Y$  and  $X$  with distributions on  $L = \{0, 1\}^4$  that satisfy top-to-bottom symmetry, the following conditions are equivalent:*

- i)  $Y \succ_{SPM} X$ ;
- ii)  $Y \succ_{CXMOD} X$ ;
- iii)  $Y \succ_{DISP} X$ ;
- iv)  $c(Y) \succ_{CX} c(X)$  and for all  $i \neq j$ ,  $Pr(Y_i = Y_j) \geq Pr(X_i = X_j)$ .

The proof, in Appendix D, has a similar structure to that of Proposition 6. We use Proposition 4 to show that when  $L = \{0, 1\}^4$  and the distributions satisfy top-to-bottom symmetry,  $Y \succ_{DISP} X$

implies that iv) holds. (Top-to-bottom symmetry on  $L = \{0, 1\}^4$  itself ensures that  $Y$  and  $X$  have identical marginal distributions, since it implies that for all  $i$ ,  $Pr(Y_i = 1) = Pr(Y_i = 0) = \frac{1}{2}$  and  $Pr(X_i = 1) = Pr(X_i = 0) = \frac{1}{2}$ .) We then adapt the construction used to prove Proposition 6 to show that the conditions in iv) ensure the existence of a sequence of nonnegative ETs that sum to  $g - f$ . Since each ET raises the expectation of any supermodular function,  $g \succ_{SPM} f$  and hence, by Theorem 2, conditions i)–iv) in Proposition 7 are all equivalent.

Even though, in the environment of Proposition 7, the SPM, CXMOD, and DISP orderings are equivalent, we can show by example that the GWA ordering is strictly stronger and the concordance ordering strictly weaker.<sup>26</sup>

## 5. Conclusion

In this paper we examined five orderings of interdependence for  $n$ -dimensional distributions and described a range of economic contexts in which these orderings are applicable, including multidimensional inequality comparisons, assessments of ex post inequality under uncertainty, valuations of portfolios of assets or insurance policies, and assessments of systemic risk in financial systems. While greater weak association, the supermodular ordering, and the concordance ordering have received some attention in the statistics and economics literatures, this paper introduces the dispersion ordering and the convex-modular ordering. Our dispersion ordering is motivated by the link between greater interdependence in random vectors and lower dispersion of the cdf's of their order statistics, and we proved that greater interdependence for  $Y$  than for  $X$  according to the dispersion ordering corresponds to greater riskiness, in the sense of Rothschild and Stiglitz [48], of the summary statistic  $\sum_{i=1}^n I_{\{Y_i > s_i\}}$  compared to  $\sum_{i=1}^n I_{\{X_i > s_i\}}$ , for any point  $(s_1, \dots, s_n)$  in the support. Similarly,  $Y$  is more interdependent than  $X$  according to the convex-modular ordering if for any choice of  $\{r_i\}_{i=1}^n$  nondecreasing, the aggregate  $\sum_{i=1}^n r_i(Y_i)$  is riskier than  $\sum_{i=1}^n r_i(X_i)$ . Theorem 1 showed that for  $n = 2$ , all five orderings of interdependence are equivalent. In contrast, Theorem 2 showed that for  $n = 3$ , four of the orderings are strictly ranked, and only the dispersion and concordance orderings are equivalent, while for  $n > 3$ , all five orderings are strictly ranked. For multivariate random vectors each of whose components are binary, we demonstrated some equivalences among the orderings and provided easily checkable and easily interpretable necessary and sufficient conditions for the orderings to hold. For arbitrary multivariate distributions, we emphasized that the dispersion and concordance orderings have the advantage, relative to the other three, of being checkable pointwise.

## Appendix A. Proof that $Y \succ_{GWA} X$ implies $Y \succ_{SPM} X$

This proof builds on Christofides and Vaggelatos's [18] proof that if  $(Y_1, \dots, Y_n)$  is weakly associated, then  $(Y_1, \dots, Y_n)$  dominates its independent counterpart according to the supermod-

<sup>26</sup> To show that the GWA ordering is strictly stronger, let  $X$  have a uniform distribution on  $\{0, 1\}^4$  and let  $Y$  be obtained from  $X$  by two ETs as defined in (4), each of size  $\alpha \in (0, \frac{1}{16})$ , one involving  $(0, 0, 1, 1)$ ,  $(1, 0, 1, 1)$ ,  $(0, 1, 1, 1)$ , and  $(1, 1, 1, 1)$  and the other involving  $(0, 0, 0, 0)$ ,  $(1, 0, 0, 0)$ ,  $(0, 1, 0, 0)$ , and  $(1, 1, 0, 0)$ . Both  $Y$  and  $X$  satisfy top-to-bottom symmetry, and by construction,  $Y \succ_{SPM} X$ . However, for  $r(z_1, z_2, z_3) = I_{\{z_1+z_2+z_3 \geq 2\}}$  and  $s(z_4) = I_{\{z_4=1\}}$ ,  $Cov(r(Y_1, Y_2, Y_3), s(Y_4)) < 0 = Cov(r(X_1, X_2, X_3), s(X_4))$ , so  $Y \not\succ_{GWA} X$  does not hold. To show that the concordance ordering is strictly weaker, recall Example 3, which was used in part iv) of the proof of Theorem 2. In that example,  $\delta = g - f$  satisfies top-to-bottom symmetry, and as we showed,  $Y \succ_{CONC} X$  holds but  $Y \not\succ_{DISP} X$  does not.



ular ordering. First, we define a new random vector  $Z$  that has the same distribution as  $X$  and is independent of  $Y$ . We then show by induction that  $Y \succ_{SPM} Z$  and hence  $Y \succ_{SPM} X$ .

For  $n = 2$ , the result is proved in the references cited in the proof of [Theorem 1](#). Suppose that it is true for  $n = m - 1$ , so that for all supermodular  $w$ ,  $Ew(Y_2, \dots, Y_m) \geqslant Ew(Z_2, \dots, Z_m)$ . Then

$$\begin{aligned}
 Ew(Z_1, Y_2, \dots, Y_m) &= \sum_{i=0}^{l_i-1} E[w(i, Y_2, \dots, Y_m) | Z_1 = i] Pr(Z_1 = i) \\
 &= \sum_{i=0}^{l_i-1} E[w(i, Y_2, \dots, Y_m)] Pr(Z_1 = i) \tag{20}
 \end{aligned}$$

$$= \sum_{i=0}^{l_i-1} E[w(i, Y_2, \dots, Y_m)] Pr(Y_1 = i) \tag{21}$$

$$\begin{aligned}
 &\geqslant \sum_{i=0}^{l_i-1} E[w(i, Z_2, \dots, Z_m)] Pr(Y_1 = i) \\
 &= \sum_{i=0}^{l_i-1} E[w(i, Z_2, \dots, Z_m) | Y_1 = i] Pr(Y_1 = i) \tag{22} \\
 &= Ew(Y_1, Z_2, \dots, Z_m).
 \end{aligned}$$

The equality in (20) uses the independence of  $Z$  from  $Y$ , and that in (21) the assumption that  $Y$  and  $X$ , hence  $Y$  and  $Z$ , have identical marginal distributions (along with the fact that  $E[w(i, Y_2, \dots, Y_m)]$  is a univariate function of  $i$ ). The inequality follows by applying the induction hypothesis for each value of  $i$ . The equality in (22) uses the independence of  $Z$  from  $Y$ .

Now we show that for all supermodular  $w$ ,

$$\begin{aligned}
 &Ew(Y_1, Y_2, \dots, Y_m) - Ew(Z_1, Z_2, \dots, Z_m) \\
 &\geqslant Ew(Z_1, Y_2, \dots, Y_m) - Ew(Y_1, Z_2, \dots, Z_m),
 \end{aligned}$$

which combined with the previous steps yields the desired result. First observe that we can write

$$\begin{aligned}
 &w(Y_1, Y_2, \dots, Y_m) - w(Z_1, Y_2, \dots, Y_m) \\
 &= \sum_{i=0}^{l_i-1} (I_{\{Y_1 > i\}} - I_{\{Z_1 > i\}}) (w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m)).
 \end{aligned}$$

Hence,

$$\begin{aligned}
 &Ew(Y_1, Y_2, \dots, Y_m) - Ew(Z_1, Y_2, \dots, Y_m) \\
 &= \sum_{i=0}^{l_i-1} \{ E[I_{\{Y_1 > i\}} (w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m))] \\
 &\quad - E[I_{\{Z_1 > i\}} (w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m))] \} \\
 &= \sum_{i=0}^{l_i-1} \{ E[I_{\{Y_1 > i\}} (w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m))] \\
 &\quad - E[I_{\{Z_1 > i\}} (w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m))] \}
 \end{aligned}$$

$$- E[I_{\{Z_1>i\}}]E[w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m)]\} \tag{23}$$

$$= \sum_{i=0}^{l_i-1} \{E[I_{\{Y_1>i\}}(w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m))]\} \\ - E[I_{\{Y_1>i\}}]E[w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m)]\} \tag{24}$$

$$= \sum_{i=0}^{l_i-1} Cov(I_{\{Y_1>i\}}, w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m)) \\ \geq \sum_{i=0}^{l_i-1} Cov(I_{\{Z_1>i\}}, w(i + 1, Z_2, \dots, Z_m) - w(i, Z_2, \dots, Z_m)) \\ = Ew(Z_1, Z_2, \dots, Z_m) - Ew(Y_1, Z_2, \dots, Z_m). \tag{25}$$

The equality in (23) uses the independence of  $Z$  from  $Y$ , and that in (24) the assumption that  $Y$  and  $X$ , hence  $Y$  and  $Z$ , have identical marginal distributions. The inequality holds since i)  $I_{\{Y_1>i\}}$  is a nondecreasing function of  $Y_1$ ; ii)  $(w(i + 1, Y_2, \dots, Y_m) - w(i, Y_2, \dots, Y_m))$  is a nondecreasing function of  $(Y_2, \dots, Y_m)$  since  $w$  is supermodular; and iii) by hypothesis,  $Y \succ_{GWA} X$  and hence  $Y \succ_{GWA} Z$ . The equality in (25) follows from the logic of the first four equalities.  $\square$

**Appendix B. Proof that for  $n \geq 3$ , the supermodular ordering is strictly stronger than the convex-modular ordering**

Since every convex-modular function is supermodular,  $Y \succ_{SPM} X$  implies  $Y \succ_{CXMOD} X$ . We show that the implication is strict by providing an example of a supermodular function on  $L = \{0, 1, 2\}^3$  which cannot be written as a nonnegative weighted sum of convex-modular ones:

Define  $w$  as follows:  $w(2, 2, 2) = 3$ ,  $w(2, 2, 1) = w(2, 1, 2) = w(1, 2, 2) = 2$ ,  $w(2, 2, 0) = w(2, 0, 2) = w(0, 2, 2) = 1$ ,  $w(2, 1, 1) = w(1, 2, 1) = w(1, 1, 2) = 1$ ,  $w(0, 2, 1) = 1$ , and  $w(z) = 0$  for all other nodes  $z \in L$ . We first show that this function  $w(x)$  is not itself convex-modular. Suppose it were. Then clearly the function  $\phi(\cdot)$  would have to take values in  $\{0, 1, 2, 3\}$ . If  $\sum_{i=1}^3 r^i(x_i)$  were strictly larger at  $(0, 2, 2)$  than at  $(0, 2, 1)$ , then since  $w(0, 2, 2) = w(0, 2, 1) = 1$ ,  $\phi(\cdot)$  would not be convex, since  $\phi(\cdot)$  would rise from 0 to 1 but then remain constant at 1 even though  $\sum_{i=1}^3 r^i(x_i)$  increased. If, instead,  $\sum_{i=1}^3 r^i(x_i)$  took on the same value at  $(0, 2, 2)$  as at  $(0, 2, 1)$ , then since  $\sum_{i=1}^3 r^i(x_i)$  is modular (additively separable) in the  $x_i$ 's, it would follow that  $\sum_{i=1}^3 r^i(x_i)$  took on the same value at  $(1, 2, 2)$  as at  $(1, 2, 1)$ . However,  $w(1, 2, 2) = 2 > 1 = w(1, 2, 1)$ . Thus, we reach a contradiction, so  $w(z)$  as defined above is not convex-modular.

In Meyer and Strulovici [42], we show how, for any finite support  $L$ , the “double description method”, conceptualized by Motzkin et al. [45], can be used to determine the extreme rays of the cone of supermodular functions on  $L$ . For  $L = \{0, 1, 2\}^3$ , we show there that the function  $w$  defined above is an extreme ray of the cone of supermodular functions and hence cannot be non-trivially expressed as a nonnegative weighted sum of supermodular functions. This, combined with the fact that it is not itself convex-modular, shows that it cannot be expressed as a nonnegative weighted sum of convex-modular functions.  $\square$




Fig. 2.  $\delta(z) = g(z) - f(z)$  for  $L = \{0, 1\}^4$  and with “top-to-bottom symmetry” (which implies that  $c_{14} = c_{23}$ ,  $c_{24} = c_{13}$ , and  $c_{34} = c_{12}$ ). In the left cube,  $z_4 = 0$ ; in the right cube,  $z_4 = 1$ .

**Appendix C. Proof that in the example in Remark 2 (Section 3),  $\tilde{g} \succ_{GWA} \tilde{f}$**

Since  $\tilde{f}$  and  $\tilde{g}$  are symmetric distributions, all partitions of  $\{Z_1, Z_2, Z_3\}$  into disjoint sets  $\{Z_i\}$  and  $\{Z_j, Z_k\}$  will yield the same covariances. We therefore focus on the partition into  $\{Z_1\}$  and  $\{Z_2, Z_3\}$ . As is well known, any nondecreasing function on  $L = \{0, 1\}^3$  can be written as a non-negative weighted combination of indicator functions, so it is sufficient to focus on functions  $r, s$  that are nondecreasing indicator functions. The only non-trivial nondecreasing indicator function of  $z_1$  is  $r(z_1) = I_{\{z_1=1\}}$ . There are four non-trivial nondecreasing indicator functions of  $(z_2, z_3)$ :  $s(z_2, z_3) = I_{\{z_2=1\}}$ ,  $s(z_2, z_3) = I_{\{z_3=1\}}$ ,  $s(z_2, z_3) = I_{\{z_2=z_3=1\}}$ , and  $s(z_2, z_3) = I_{\{z_2+z_3 \geq 1\}}$ .

First take  $s(z_2, z_3) = I_{\{z_2=1\}}$ . Recall that  $\tilde{g} - \tilde{f} = g - f$ , where  $g$  and  $f$  are defined in Example 1. Therefore, since it was shown in Example 1 that  $g \succ_{SPM} f$  and since the supermodular ordering is a difference-based order,  $\tilde{g} \succ_{SPM} \tilde{f}$ . Note also that  $\tilde{g}$  and  $\tilde{f}$  have identical marginal distributions, so  $Er(Y_1) = Pr(Y_1 = 1) = Pr(X_1 = 1) = Er(X_1)$  and  $Es(Y_2, Y_3) = Pr(Y_2 = 1) = Pr(X_2 = 1) = Es(X_2, X_3)$ . Therefore,  $Cov(r(Y_1), s(Y_2, Y_3)) - Cov(r(X_1), s(X_2, X_3)) = E[r(Y_1)s(Y_2, Y_3)] - E[r(X_1)s(X_2, X_3)]$ . Since  $r(z_1)s(z_2, z_3) = I_{\{z_1=z_2=1\}}$  is supermodular, it thus follows from  $\tilde{g} \succ_{SPM} \tilde{f}$  that  $Cov(r(Y_1), s(Y_2, Y_3)) \geq Cov(r(X_1), s(X_2, X_3))$ . An analogous argument holds if  $s(z_2, z_3) = I_{\{z_3=1\}}$ .

It remains only to consider  $s(z_2, z_3) = I_{\{z_2=z_3=1\}}$  and  $s(z_2, z_3) = I_{\{z_2+z_3 \geq 1\}}$ . For  $s(z_2, z_3) = I_{\{z_2=z_3=1\}}$ ,  $Cov(r(Y_1), s(Y_2, Y_3)) - Cov(r(X_1), s(X_2, X_3)) = 0$ , while for  $s(z_2, z_3) = I_{\{z_2+z_3 \geq 1\}}$ ,  $Cov(r(Y_1), s(Y_2, Y_3)) > 0 > Cov(r(X_1), s(X_2, X_3))$ . Therefore,  $\tilde{g} \succ_{GWA} \tilde{f}$ .  $\square$

**Appendix D. Proof of Proposition 7**

Since the SPM, CXMOD, DISP, and CX orderings are all difference-based orderings, it is sufficient to work with  $\delta \equiv g - f$ . Fig. 2 displays the values of  $\delta$  at each of the 16 nodes of  $L = \{0, 1\}^4$ , given top-to-bottom symmetry of  $g$  and  $f$ , and hence of  $\delta$ . Top-to-bottom symmetry of  $\delta$  implies that once the 8 values of  $\delta(1, 1, 1, 1) \equiv a$ ,  $\delta(0, 1, 1, 1) \equiv b_1$ ,  $\delta(1, 0, 1, 1) \equiv b_2$ ,  $\delta(1, 1, 0, 1) \equiv b_3$ ,  $\delta(1, 1, 1, 0) \equiv b_4$ ,  $\delta(0, 1, 1, 0) \equiv c_{14}$ ,  $\delta(1, 0, 1, 0) \equiv c_{24}$ , and  $\delta(1, 1, 0, 0) \equiv c_{34}$  are specified (corresponding to the 8 nodes marked with black dots in Fig. 2), the remaining 8 values are determined. Note that top-to-bottom symmetry implies that  $c_{12} = c_{34}$ ,  $c_{13} = c_{24}$ ,

and  $c_{23} = c_{14}$ . Top-to-bottom symmetry of  $g$  and  $f$  also implies that  $Y$  and  $X$  have identical marginal distributions, since it implies that for all  $i$ ,  $Pr(Y_i = 1) = Pr(Y_i = 0) = \frac{1}{2}$  and  $Pr(X_i = 1) = Pr(X_i = 0) = \frac{1}{2}$ . From Fig. 2, identical marginals for  $Y$  and  $X$  here corresponds to

$$a + \sum_{h=1}^4 b_h + c_{ij} + c_{ik} + c_{il} = 0 \quad \forall i \neq j \neq k \neq l. \tag{26}$$

We first show that condition iii) in the proposition implies condition iv). First note that from Fig. 2, it is apparent that the second part of condition iv),  $Pr(Y_i = Y_j) \geq Pr(X_i = X_j)$  for all  $i \neq j$ , holds if and only if  $a + b_i + b_j + c_{ij} \geq 0$  for all  $i \neq j$ . Now use Proposition 4 to identify the implications of  $Y \succ_{DISP} X$ . For  $s = (0, 0, 0, 0)$ ,  $Y_i^s = Y_i$  and  $X_i^s = X_i$  for all  $i$ , so  $c^s(Y^s) \succ_{CX} c^s(X^s)$  if and only if  $c(Y) \succ_{CX} c(X)$ . For  $s = (1, 1, 0, 0)$  and all permutations thereof,  $c^s(Y^s)$  and  $c^s(X^s)$  have support  $\{0, 1, 2\}$  and, from Eq. (15),  $c^s(Y^s) \succ_{CX} c^s(X^s)$  implies that  $E[\max\{c^s(Y^s) - 1, 0\}] - E[\max\{c^s(X^s) - 1, 0\}] \geq 0$ , which is equivalent to  $a + b_i + b_j + c_{ij} \geq 0$  for all  $i \neq j$ . Hence iii) implies iv).

In order to show that condition iv) implies condition i), we identify the inequalities to which  $c(Y) \succ_{CX} c(X)$  is equivalent here. From (15),  $c(Y) \succ_{CX} c(X)$  if and only if 1)  $Ec(Y) - Ec(X) = 0$  and 2)  $E[\max\{c(Y) - t, 0\}] - E[\max\{c(X) - t, 0\}] \geq 0$  for  $t = 1, 2, 3$ . Given top-to-bottom symmetry,  $Ec(Y) - Ec(X) = 4a + 3(\sum_{h=1}^4 b_h) + 2 \cdot 2(c_{ij} + c_{ik} + c_{il}) + \sum_{h=1}^4 b_h$ , which equals 0 given (26). For  $t = 3$ , condition 2) corresponds to  $a \geq 0$ , and for  $t = 2$ , it corresponds to  $2a + \sum_{h=1}^4 b_h \geq 0$ . For  $t = 1$  and given top-to-bottom symmetry, the left-hand side of condition 2) is  $3a + 2 \sum_{h=1}^4 b_h + 2(c_{ij} + c_{ik} + c_{il})$ , which equals  $a$  given (26). Therefore,  $c(Y) \succ_{CX} c(X)$  here if and only if  $a \geq 0$  and  $2a + \sum_{h=1}^4 b_h \geq 0$ .

We now show that

$$a \geq 0, \quad 2a + \sum_{h=1}^4 b_h \geq 0, \quad \text{and} \quad \forall i \neq j \quad a + b_i + b_j + c_{ij} \geq 0 \quad \Rightarrow \quad g \succ_{SPM} f.$$

To do so, we decompose  $\delta$  into 24 elementary transformations (ETs) of the form defined in (4): for each of the 24 faces of the hypercube  $L = \{0, 1\}^4$ , there is one ET involving the 4 nodes on that face. We will abuse notation slightly and let the value of  $\delta$  at a given node also serve as the label for that node. Let the two ETs involving the nodes  $a, b_i, b_j$ , and  $c_{ij}$  (there are two such ETs because of the top-to-bottom symmetry of  $\delta$ ) have size  $\beta_{ij} = \beta_{ji}$ . Let the two ETs involving the nodes  $b_i, c_{ik}, c_{il}, b_j$  (once again, there are two such ETs because of the top-to-bottom symmetry) have size  $\alpha_{ij} = \alpha_{ji}$ . There are 6 distinct values of  $\beta_{ij}$  and 6 distinct values of  $\alpha_{ij}$ . For the 24 ETs so defined to sum to  $\delta$ , it is necessary and sufficient that  $\beta_{ij}$  and  $\alpha_{ij}$  satisfy<sup>27</sup>

$$\begin{aligned} \sum_{i < j} \beta_{ij} &= a, \\ b_i + \beta_{ij} + \beta_{ik} + \beta_{il} &= \alpha_{ij} + \alpha_{ik} + \alpha_{il} \quad \forall i \neq j \neq k \neq l, \\ -c_{ij} + \beta_{ij} + \beta_{kl} &= \alpha_{ik} + \alpha_{il} + \alpha_{jk} + \alpha_{jl} \quad \forall i \neq j \neq k \neq l. \end{aligned} \tag{27}$$

<sup>27</sup> Further details of this and subsequent arguments are provided in the supplementary material, Meyer and Strulovici [43], available on the journal website.

These three (sets of) equations ensure that each of the (sets of) nodes labeled  $a$ ,  $b_i$ , and  $c_{ij}$ , respectively, in Fig. 2 is transformed from its values under the distribution  $f$  to its values under the distribution  $g$  by the sequence of ETs just defined. Eqs. (27) can be rearranged to

$$\begin{aligned} \sum_{i < j} \beta_{ij} &= a, \\ b_i + \beta_{ij} + \beta_{ik} + \beta_{il} &= \alpha_{ij} + \alpha_{ik} + \alpha_{il} \quad \forall i \neq j \neq k \neq l, \\ 2\alpha_{ij} + 2\beta_{kl} &= a + b_i + b_j + c_{ij} \quad \forall i \neq j \neq k \neq l. \end{aligned} \tag{28}$$

Noting the similarity between Eqs. (28) and (18) for the 3-dimensional cube, set

$$\begin{aligned} \beta_{kl} &= \frac{a(a + b_i + b_j + c_{ij})}{4a + \sum_{h=1}^4 b_h}, \quad \forall i \neq j \neq k \neq l, \\ \alpha_{ij} &= \frac{(2a + \sum_{h=1}^4 b_h)(a + b_i + b_j + c_{ij})}{2(4a + \sum_{h=1}^4 b_h)}, \quad \forall i \neq j. \end{aligned} \tag{29}$$

Recalling that top-to-bottom symmetry of  $\delta$  implies that  $c_{12} = c_{34}$ ,  $c_{13} = c_{24}$ , and  $c_{23} = c_{14}$  and using (26), it can be checked, with some algebra, that with these choices for  $\beta_{kl}$  and  $\alpha_{ij}$ , Eqs. (28) are satisfied. Furthermore, if  $a \geq 0$ ,  $2a + \sum_{h=1}^4 b_h \geq 0$  and for all  $i \neq j$ ,  $a + b_i + b_j + c_{ij} \geq 0$ , then  $\beta_{kl}$  and  $\alpha_{ij}$  as defined in (29) are all nonnegative. Thus, condition iv) in the proposition implies the existence of a sequence of nonnegative ETs that sum to  $\delta = g - f$ . Since each ET raises the expectation of any supermodular function,  $g \succ_{SPM} f$ . By Theorem 2, it then follows that conditions i)–iv) are all equivalent.  $\square$

## References

- [1] V.V. Acharya, A theory of systemic risk and design of prudential bank regulation, *J. Finan. Stability* 5 (2009) 224–255.
- [2] M.D. Adler, C.W. Sanchirico, Inequality and uncertainty: theory and legal applications, *Univ. of Pennsylvania Law Rev.* 155 (2006) 279–377.
- [3] F. Allen, A. Babus, E. Carletti, Asset commonality, debt maturity and systemic risk, *J. Finan. Econ.*, doi:10.1016/j.fineco.2011.07.003, forthcoming.
- [4] A.B. Atkinson, On the measurement of inequality, *J. Econ. Theory* 2 (1970) 244–263.
- [5] A.B. Atkinson, Multidimensional deprivation: contrasting social welfare and counting approaches, *J. Econ. Inequality* 1 (2003) 51–65.
- [6] A.B. Atkinson, F. Bourguignon, The comparison of multi-dimensioned distributions of economic status, *Rev. Econ. Stud.* 49 (1982) 183–201.
- [7] K.A. Baldiga, J.R. Green, Assent-maximizing social choice, *Soc. Choice Welfare*, forthcoming.
- [8] N. Beale, D. Rand, H. Battey, K. Croxson, R. May, M. Nowak, Individual versus systemic risk and the regulator's dilemma, *Proc. Natl. Acad. Sci. USA* 31 (2011) 12647–12652.
- [9] E. Ben-Porath, I. Gilboa, D. Schmeidler, On the measurement of inequality under uncertainty, *J. Econ. Theory* 75 (1997) 194–204.
- [10] P.J. Boland, F. Proschan, Multivariate arrangement increasing functions with applications in probability and statistics, *J. Multivariate Anal.* 25 (1988) 286–298.
- [11] F. Bourguignon, S.R. Chakravarty, The measurement of multidimensional poverty, *J. Econ. Inequality* 1 (2003) 25–49.
- [12] R.M. Burton, A.R. Dabrowski, H. Dehling, An invariance principle for weakly associated random variables, *Stoch. Process. Appl.* 23 (1986) 301–306.
- [13] B. Caillaud, J. Tirole, Consensus building: how to persuade a group, *Amer. Econ. Rev.* 97 (2007) 1877–1900.
- [14] A. Calvo-Armengol, M.O. Jackson, The effects of social networks on employment and inequality, *Amer. Econ. Rev.* 94 (2004) 426–454.

- [15] A. Calvo-Armengol, M.O. Jackson, Networks in labor markets: wage and employment dynamics and inequality, *J. Econ. Theory* 132 (2007) 27–46.
- [16] L.I. de Castro, Affiliation and dependence in economic models, Discussion Paper No. 1479, Ctr. for Math. Stud. in Econ. and Mgt. Sci., Northwestern Univ., 2009.
- [17] S.H. Chew, J.S. Sagi, An inequality measure for stochastic allocations, *J. Econ. Theory* 147 (4) (2012) 1517–1544, in this issue.
- [18] T.C. Christofides, E. Vaggelatos, A connection between supermodular ordering and positive/negative association, *J. Multivariate Anal.* 88 (2004) 138–151.
- [19] S. Choi, D. Gale, S. Kariv, Behavioral aspects of learning in social networks: an experimental study, in: J. Morgan (Ed.), *Advances in Applied Microeconomics*, vol. 13: Experimental and Behavioral Economics, Emerald, Bingley, UK, 2005, pp. 25–61.
- [20] P. Dasgupta, A.K. Sen, D. Starrett, Notes on the measurement of inequality, *J. Econ. Theory* 6 (1973) 180–187.
- [21] M. Denuit, J. Dhaene, M.J. Goovaerts, R. Kaas, *Actuarial Theory for Dependent Risks: Measures, Orders, and Models*, Wiley, Chichester, West Sussex, 2005.
- [22] K. Decancq, Elementary multivariate rearrangements and stochastic dominance on a Fréchet class, *J. Econ. Theory* 147 (4) (2012) 1450–1459, in this issue.
- [23] L. Epstein, S. Tanny, Increasing generalized correlation: a definition and some economic consequences, *Can. J. Econ.* 13 (1980) 16–34.
- [24] J.D. Esary, F. Proschan, D.W. Walkup, Association of random variables, with applications, *Ann. of Math. Stat.* 44 (1967) 1466–1474.
- [25] R. Fernandez, J. Gali, To each according to ...? markets, tournaments, and the matching problem with borrowing constraints, *Rev. Econ. Stud.* 66 (1999) 799–824.
- [26] T. Gajdos, E. Maurin, Unequal uncertainties and uncertain inequalities: an axiomatic approach, *J. Econ. Theory* 116 (2004) 93–118.
- [27] A. Galeotti, S. Goyal, M.O. Jackson, F. Vega-Redondo, L. Yariv, Network games, *Rev. Econ. Stud.* 77 (2010) 218–244.
- [28] G.H. Hardy, J.E. Littlewood, G. Pólya, Some simple inequalities satisfied by convex functions, *Messenger of Maths.* 58 (1929) 145–152.
- [29] G.H. Hardy, J.E. Littlewood, G. Pólya, *Inequalities*, 2nd ed., Cambridge University Press, Cambridge, 1952.
- [30] D.A. Hennessy, H.E. Lapan, A definition of “more systematic risk” with some welfare implications, *Economica* 70 (2003) 493–507.
- [31] T. Hu, A. Müller, M. Scarsini, Some counterexamples in positive dependence, *J. Stat. Plann. Inference* 124 (2004) 153–158.
- [32] T. Hu, C. Xie, L. Ruan, Dependence structures of multivariate Bernoulli random vectors, *J. Multivariate Anal.* 94 (2005) 172–195.
- [33] I. Jewitt, Risk aversion and the choice between risky prospects: the preservation of comparative statics results, *Rev. Econ. Stud.* 54 (1987) 73–86.
- [34] K. Joag-Dev, F. Proschan, Negative association of random variables, with applications, *Ann. of Stat.* 11 (1983) 286–295.
- [35] H. Joe, Multivariate concordance, *J. Multivariate Anal.* 35 (1990) 12–30.
- [36] H. Joe, *Multivariate Models and Dependence Concepts*, Chapman and Hall/CRC, London, 1997.
- [37] H. Levy, J. Paroush, Toward multivariate efficiency criteria, *J. Econ. Theory* 7 (1974) 129–142.
- [38] R.P. McAfee, Coarse matching, *Econometrica* 70 (2002) 2025–2034.
- [39] M.A. Meyer, Interdependence in multivariate distributions: stochastic dominance theorems and an application to the measurement of ex post inequality under uncertainty, Nuffield Coll. Discussion Paper No. 49, 1990.
- [40] M.A. Meyer, D. Mookherjee, Incentives, compensation and social welfare, *Rev. Econ. Stud.* 45 (1987) 209–226.
- [41] M.A. Meyer, M. Rothschild, (In)efficiency in sorting and matching, unpublished manuscript, Oxford, 2004.
- [42] M.A. Meyer, B. Strulovici, The supermodular stochastic ordering, unpublished manuscript, Oxford and Northwestern, 2011.
- [43] M.A. Meyer, B. Strulovici, Supplementary material for “Increasing Interdependence of Multivariate Distributions”, 2011.
- [44] P.R. Milgrom, R.J. Weber, A theory of auctions and competitive bidding, *Econometrica* 50 (1982) 1089–1122.
- [45] T.S. Motzkin, H. Raiffa, G.L. Thompson, R.M. Thrall, The double description method, in: H.W. Kuhn, A.W. Tucker (Eds.), *Contributions to the Theory of Games II*, Princeton University Press, Princeton, 1953, pp. 51–73.
- [46] A. Müller, D. Stoyan, *Comparison Methods for Stochastic Models and Risks*, Wiley, Chichester, West Sussex, 2002.
- [47] A. Prat, Should a team be homogeneous?, *Europ. Econ. Rev.* 46 (2002) 1187–1207.

- [48] M. Rothschild, J.E. Stiglitz, Increasing risk: I. A definition, *J. Econ. Theory* 2 (1970) 225–243.
- [49] M. Shaked, J.G. Shanthikumar, Supermodular stochastic orders and positive dependence of random vectors, *J. Multivariate Anal.* 61 (1997) 86–101.
- [50] M. Shaked, Y.L. Tong, Some partial orderings of exchangeable random variables by positive dependence, *J. Multivariate Anal.* 17 (1985) 333–349.
- [51] R. Shimer, L. Smith, Assortative matching and search, *Econometrica* 68 (2000) 343–369.
- [52] A.H. Tchen, Inequalities for distributions with given marginals, *Ann. of Prob.* 8 (1980) 814–827.
- [53] D.M. Topkis, Minimizing a submodular function on a lattice, *Oper. Res.* 26 (1978) 305–331.