

Dynamic Factor Models

January 2010
This revision: May 7, 2010

James H. Stock

Department of Economics, Harvard University
and the National Bureau of Economic Research

and

Mark W. Watson*

Woodrow Wilson School and Department of Economics, Princeton University
and the National Bureau of Economic Research

*Prepared for the *Oxford Handbook of Economic Forecasting*, Michael P. Clements and David F. Hendry (eds), Oxford University Press. We thank Jushan Bai and Serena Ng for helpful discussions and Ugo Troiano for research assistance.

1. Introduction

Macroeconometricians face a peculiar data structure. On the one hand, the number of years for which there is reliable and relevant data is limited and cannot readily be increased other than by the passage of time. On the other hand, for much of the postwar period statistical agencies have collected monthly or quarterly data on a great many related macroeconomic, financial, and sectoral variables. Thus macroeconometricians face data sets that have hundreds or even thousands of series, but the number of observations on each series is relatively short, for example 20 to 40 years of quarterly data.

This chapter surveys work on a class of models, dynamic factor models (DFMs), which has received considerable attention in the past decade because of their ability to model simultaneously and consistently data sets in which the number of series exceeds the number of time series observations. Dynamic factor models were originally proposed by Geweke (1977) as a time-series extension of factor models previously developed for cross-sectional data. In early influential work, Sargent and Sims (1977) showed that two dynamic factors could explain a large fraction of the variance of important U.S. quarterly macroeconomic variables, including output, employment, and prices. This central empirical finding that a few factors can explain a large fraction of the variance of many macroeconomic series has been confirmed by many studies; see for example Giannone, Reichlin, and Sala (2004) and Watson (2004).

The aim of this survey is to describe, at a level that is specific enough to be useful to researchers new to the area, the key theoretical results, applications, and empirical findings in the recent literature on DFMs. Bai and Ng (2008) and Stock and Watson

(2006) provide complementary surveys of this literature. Bai and Ng's (2008) survey is more technical than this one and focuses on the econometric theory and conditions; Stock and Watson (2006) focus on DFM-based forecasts in the context of other methods for forecasting with many predictors.

The premise of a dynamic factor model is that a few latent dynamic factors, f_t , drive the comovements of a high-dimensional vector of time-series variables, X_t , which is also affected by a vector of mean-zero idiosyncratic disturbances, e_t . These idiosyncratic disturbances arise from measurement error and from special features that are specific to an individual series (the effect of a Salmonella scare on restaurant employment, for example). The latent factors follow a time series process, which is commonly taken to be a vector autoregression (VAR). In equations, the dynamic factor model is,

$$X_t = \lambda(L)f_t + e_t \quad (1)$$

$$f_t = \Psi(L)f_{t-1} + \eta_t \quad (2)$$

where there are N series, so X_t and e_t are $N \times 1$, there are q dynamic factors so f_t and η_t are $q \times 1$, L is the lag operator, and the lag polynomial matrices $\lambda(L)$ and $\Psi(L)$ are $N \times q$ and $q \times q$, respectively. The i^{th} lag polynomial $\lambda_i(L)$ is called the dynamic factor loading for the i^{th} series, X_{it} , and $\lambda_i(L)f_t$ is called the common component of the i^{th} series. We assume that all the processes in (1) and (2) are stationary (nonstationarity is discussed in the final section of this chapter). The idiosyncratic disturbances are assumed to be uncorrelated with the factor innovations at all leads and lags, that is, $Ee_t\eta_{t-k}' = 0$ for all k . In the so-called exact dynamic factor model, the idiosyncratic disturbances are assumed to be mutually uncorrelated at all leads and lags, that is, $Ee_{it}e_{js} = 0$ for all s if $i \neq j$.

An important motivation for considering DFMs is that, if one knew the factors f_t and if (e_t, η_t) are Gaussian, then one can make efficient forecasts for an individual variable using the population regression of that variable on the lagged factors and lags of that variable. Thus the forecaster gets the benefit of using all N variables by using only q factors, where q is typically much smaller than N . Specifically, under squared error loss, the optimal one-step ahead forecast of the i^{th} variable is,

$$\begin{aligned}
E[X_{it+1} | X_t, f_t, X_{t-1}, f_{t-1}, \dots] &= E[\lambda_i(L)f_{t+1} + e_{it+1} | X_t, f_t, X_{t-1}, f_{t-1}, \dots] \\
&= E[\lambda_i(L)f_{t+1} | X_t, f_t, X_{t-1}, f_{t-1}, \dots] + E[e_{it+1} | X_t, f_t, X_{t-1}, f_{t-1}, \dots] \\
&= E[\lambda_i(L)f_{t+1} | f_t, f_{t-1}, \dots] + E[e_{it+1} | e_{it}, e_{it-1}, \dots] \\
&= \alpha(L)f_t + \delta(L)X_{it}, \tag{3}
\end{aligned}$$

where the third line follows from (2) and the final line follows from (1) and the exact DFM assumptions. Thus the dimension of the efficient population forecasting regression does not increase as one adds variables to the system.

The first issue at hand for the econometrician is to estimate the factors (or, more precisely, to estimate the space spanned by the factors) and to ascertain how many factors there are; these two topics are covered in the Sections 2 and 3 of this survey. Once one has reliable estimates of the factors in hand, there are a number of things one can do with them beyond using them for forecasting, including using them as instrumental variables, estimating factor-augmented vector autoregressions (FAVARs), and estimating dynamic stochastic general equilibrium models (DSGEs); these applications are covered in Section 4. Section 5 discusses some extensions.

2. Factor Estimation

The seminal work of Geweke (1977) and Sargent and Sims (1977) used frequency domain methods to look for evidence of a dynamic factor structure and to estimate the importance of the factor. Those methods, however, could not estimate f_i directly and thus could not be used for forecasting. Therefore subsequent work on DFMs focused on time-domain methods in which f_i could be estimated directly.

Work on time-domain estimation of DFMs can be divided into three generations. The first generation consisted of low-dimensional (small N) parametric models estimated in the time domain using Gaussian maximum likelihood estimation (MLE) and the Kalman filter. This method provides optimal estimates of f (and optimal forecasts) under the model assumptions and parameters. However, estimation of those parameters entails nonlinear optimization, which historically had the effect of restricting the number of parameters, and thus the number of series, that could be handled. The second generation of estimators entailed nonparametric estimation with large N using cross-sectional averaging methods, primarily principal components and related methods. The key result in this second generation is that the principal components estimator of the space spanned by the factors is consistent and moreover, if N is sufficiently large, then the factors are estimated precisely enough to be treated as data in subsequent regressions. The third generation uses these consistent nonparametric estimates of the factors to estimate the parameters of the state space model used in the first generation and thereby solves the dimensionality problem associated encountered by first-generation models. A different solution to the problem of the very many unknown parameters in the state space model is to use Bayesian methods, that is, to specify a prior and integrate instead of maximize, and

a small number of papers took this approach contemporaneously with the work on the second and third generation (classical) estimators.

The expressions in this chapter omit intercepts, and all the methods assume that the data have been transformed to eliminate unit roots and trends. Typically this is accomplished by differencing series as needed, then by standardizing the differenced series; for example, a typical element of X_{it} might be the one-period growth rate of a real activity indicator, standardized to have mean zero and unit standard deviation.

2.1 First generation: time-domain maximum likelihood via the Kalman filter

Early time-domain estimation of dynamic factor models used the Kalman filter to compute the Gaussian likelihood, estimated the parameters by maximum likelihood, then used the Kalman filter and smoother to obtain efficient estimates of the factors (Engle and Watson (1981,1983), Stock and Watson (1989), Sargent (1989), and Quah and Sargent (1993)). The first step in implementing this approach is to write the DFM as a linear state space model. Let p be the degree of the lag polynomial matrix $\lambda(L)$, let $F_t = (f_t', f_{t-1}', \dots, f_{t-p}')$ denote an $r \times 1$ vector, and let $\Lambda = (\lambda_0, \lambda_1, \dots, \lambda_p)$, where λ_i is the $N \times q$ matrix of coefficients on the i^{th} lag in $\lambda(L)$. Similarly, let $\Phi(L)$ be the matrix consisting of 1's, 0's, and the elements of $\Psi(L)$ such that the vector autoregression in (2) is rewritten in terms of F_t . With this notation the DFM (1) and (2) can be rewritten,

$$X_t = \Lambda F_t + e_t \quad (4)$$

$$\Phi(L)F_t = G\eta_t, \quad (5)$$

where G is a matrix of 1's and 0's chosen so that (5) and (2) are equivalent. Equations (4) and (5) are referred to as the “static form” of the DFM because the factors appear to enter only contemporaneously, although this is just a notational artifact since the static

factors F_t contain current and past values of the dynamic factors f_t . The linear state space model is completed by specifying a process for e_t and for the errors η_t . Typically the errors e_t are assumed to follow univariate autoregressions,

$$d_i(L)e_t = \zeta_{it}, i = 1, \dots, N. \quad (6)$$

With the further assumptions that ζ_{it} is i.i.d. $N(0, \sigma_{\zeta_i}^2)$, $i = 1, \dots, N$, η_{jt} is i.i.d. $N(0, \sigma_{\eta_j}^2)$, $j = 1, \dots, q$, and $\{\zeta_{it}\}$ and $\{\eta_{jt}\}$ are independent, equations (4) – (6) constitute a complete linear state space model. Given the parameters, the Kalman filter can be used to compute the likelihood and to estimate filtered values of F_t and thus of f_t .

An advantage of this parametric state space formulation is that it can handle data irregularities. For example, if some series are observed weekly and some are observed monthly, the latent process for the factor (5) can be formulated as evolving on a weekly time scale, but the dimension of the measurement equation (4) depends on which series are actually observed, that is, the row dimension of Λ would change depending on the variables actually observed at a given date; see Harvey (1989, p. 325) for a general discussion. Angelini, Bańbura, and Rünstler (2008) implement a DFM-based model with mixed monthly and quarterly data for the purpose of monthly distribution of Euro-area GDP, that is, for estimating and forecasting unobserved monthly GDP. In a closely related application to U.S. data, Aruoba, Diebold, and Scotti (2009) implement a DFM with a single dynamic factor and a weekly variable, four monthly variables, and a quarterly variable to produce an index of economic activity that can be updated weekly. If some series are available for only a subset of the sample, the dimension of the measurement equation can change as time series become available.

The EM algorithm can be used to compute the MLEs of the parameters.

Nevertheless, the number of parameters is proportional to N , so direct estimation of the coefficients by MLE is cumbersome and historically was prohibitive for large systems.

2.2 Second generation: nonparametric averaging methods

Why cross-sectional averaging works. The motivation for considering factor estimation by cross-sectional averaging of X_t is that weighted averages of the idiosyncratic disturbances will converge to zero by the weak law of large numbers, so that only linear combinations of the factors remain. The cross-sectional averaging estimators are based on the static representation of the DFM (4).

The cross-sectional averaging estimators are nonparametric, in the sense that they do not require a parametric model for the factors F_t as in (5) or for the idiosyncratic dynamics as in (6). Instead, F_t is treated as a r -dimensional parameter to be estimated using a N -dimensional data vector X_t . Instead of parametric assumptions, weaker assumptions along the lines of Chamberlain and Rothschild's (1983) approximate factor model are made about the factor structure. In particular, consider the conditions,

$$N^{-1}\Lambda'\Lambda \rightarrow D_\Lambda, \text{ where the } r \times r \text{ matrix } D_\Lambda \text{ has full rank, and} \quad (7)$$

$$\max_{\text{eval}}(\Sigma_e) \leq c < \infty \text{ for all } N, \quad (8)$$

where \max_{eval} denotes the maximum eigenvalue, $\Sigma_e = Ee_t e_t'$, and the limit in (7) is taken as $N \rightarrow \infty$. Condition (7) assures that the factors are pervasive (they affect most or all of the series) and that the factor loadings are heterogeneous (so that the columns of Λ are not too similar). Condition (8) ensures that the idiosyncratic disturbances have limited correlation across series.

To develop intuition for why cross-sectional averaging works, consider the estimator of F_t constructed as the weighted average of X_t using a nonrandom $N \times r$ matrix of weights W , where W is normalized so that $W'W/N = I_r$:

$$\hat{F}_t(N^{-1}W) = N^{-1}W'X_t. \quad (9)$$

If $N^{-1}W'\Lambda \rightarrow H$ as $N \rightarrow \infty$ where the $r \times r$ matrix H has full rank, and if conditions (7) and (8) hold, then $\hat{F}_t(N^{-1}W)$ is consistent for the space spanned by F_t :

$$\hat{F}_t(N^{-1}W) = N^{-1}W'(\Lambda F_t + e_t) = N^{-1}W'\Lambda F_t + N^{-1}W'e_t \xrightarrow{p} HF_t \text{ as } N \rightarrow \infty \quad (10)$$

because $N^{-1}W'\Lambda \rightarrow H$ by assumption and because $N^{-1}W'e_t \xrightarrow{p} 0$ by the weak law of large numbers.¹ Because H is full rank, $\hat{F}_t(N^{-1}W)$ consistently estimates the space spanned by the factors.

The key to the consistency result (10) is having a weight matrix W such that $N^{-1}W'\Lambda \rightarrow H$. In some special cases it is possible to use postulated structure of the DFM to construct such a W . An example that draws on Forni and Reichlin (1998) is the estimation of a single factor ($r=1$) that enters only contemporaneously in conceptually related series such as employment by sector. In this case, F_t is a scalar (in terms of (1), $\lambda_i(L)$ has degree zero and $q=1$). Further suppose that $0 < \Lambda_{min} \leq \Lambda_i \leq \Lambda_{max} < \infty$ for all i and N (this assumption implies (7)) and that (8) holds. Then $W = \iota$ (where ι is the vector of 1's) satisfies both the normalization condition $N^{-1}W'W = 1$ and the spanning condition $N^{-1}W'\Lambda \rightarrow \bar{\Lambda} > 0$, so (10) holds. Note that there are many W 's that produce a consistent

¹ The result $N^{-1}W'e_t \xrightarrow{p} 0$ follows from Chebyshev's inequality. Let W_j be the j^{th} column of W . Then $\text{var}(N^{-1}W_j'e_t) \leq \text{maxeval}(\Sigma_e)/N \leq c/N \rightarrow 0$, where the first inequality follows because W is normalized so that $N^{-1}W'W = I_r$ and the second inequality follows from (8).

estimator of F_t in this example, including random W 's. For example, $W_i = (1 + z_i)/\sqrt{2}$, where z_i is i.i.d. $N(0,1)$, satisfies $N^{-1}W'W \xrightarrow{p} 1$ and $N^{-1}W'\Lambda \xrightarrow{p} \bar{\Lambda}/\sqrt{2} > 0$, so that (10) holds with $H = \bar{\Lambda}/\sqrt{2}$. Thus many different cross sectional weighting schemes can yield a consistent estimator of F_t .

In general, there will be insufficient structure on Λ to posit a weight matrix W that does not depend on the data, which is where principal components analysis comes in.

Principal components estimation. The principal components estimator of F_t is the weighted averaging estimator (9), with $W = \hat{\Lambda}$, where $\hat{\Lambda}$ is the matrix of eigenvectors of the sample variance matrix of X_t , $\hat{\Sigma}_X = T^{-1} \sum_{t=1}^T X_t X_t'$, associated with the r largest eigenvalues of $\hat{\Sigma}_X$. The principal components estimator can be derived as the solution to the least squares problem,

$$\min_{F_1, \dots, F_T, \Lambda} V_r(\Lambda, F), \text{ where } V_r(\Lambda, F) = \frac{1}{NT} \sum_{t=1}^T (X_t - \Lambda F_t)'(X_t - \Lambda F_t), \quad (11)$$

subject to the normalization $N^{-1}\Lambda'\Lambda = I_r$. To solve (11), first minimize over F_t given Λ to

obtain $\hat{F}_t(\Lambda(\Lambda'\Lambda)^{-1}) = (\Lambda'\Lambda)^{-1} \Lambda'X_t$, then concentrate the objective function so that

(11) becomes $\min_{\Lambda} T^{-1} \sum_{t=1}^T X_t' [I - \Lambda(\Lambda'\Lambda)^{-1} \Lambda] X_t$. This minimization problem is

equivalent to $\max_{\Lambda} \text{tr} \{ (\Lambda'\Lambda)^{-1/2} \Lambda' (T^{-1} \sum_{t=1}^T X_t X_t') \Lambda (\Lambda'\Lambda)^{-1/2} \}$, which is in turn

equivalent to $\max_{\Lambda} \Lambda' \hat{\Sigma}_{XX} \Lambda$ subject to $N^{-1}\Lambda'\Lambda = I_r$. The solution to this final problem is

to set $\hat{\Lambda}$ equal to the scaled eigenvectors of $\hat{\Sigma}_{XX}$ corresponding to its r largest

eigenvalues. Because $\hat{\Lambda}'\hat{\Lambda} = NI_r$, it follows that the least squares estimator of F_t is $\hat{F}_t = \hat{F}_t(N^{-1}\hat{\Lambda}) = N^{-1}\hat{\Lambda}'X_t$, which are the scaled first r principal components of X_t .

Consistency of the principal components estimator of F_t was first shown for T fixed and $N \rightarrow \infty$ in the exact static factor model by Connor and Korajczyk (1986). Stock and Watson (2002a) proved uniform consistency of the factors under weaker conditions along the lines of Chamberlain and Rothschild's (1983) approximate factor model, allowing for weak serial and cross-correlation in the idiosyncratic errors. Stock and Watson (2002a) also provided rate conditions on N and T under which \hat{F}_t can be treated as data for the purposes of a second stage least squares regression (that is, in which the estimation error in \hat{F}_t does not affect the asymptotic distribution of the OLS coefficients with \hat{F}_t as a regressor). Bai (2003) provides limiting distributions for the estimated factors and common components. Bai and Ng (2006a) provide improved rates, specifically $N \rightarrow \infty$, $T \rightarrow \infty$, and $N^2/T \rightarrow \infty$, under which \hat{F}_t is consistent and can be treated as data in subsequent regressions; they also provide results for construction of confidence intervals for common components estimated using \hat{F}_t .

Generalized principal components estimation. Generalized principal components is to principal components as generalized least squares is to least squares. If the idiosyncratic error variance matrix Σ_e is not proportional to the identity matrix, then the analogy to least squares regression suggests that F_t and Λ solve a weighted version of (11), where the weighting matrix is Σ_e^{-1} :

$$\min_{F_1, \dots, F_T, \Lambda} T^{-1} \sum_{t=1}^T (X_t - \Lambda F_t)' \Sigma_e^{-1} (X_t - \Lambda F_t). \quad (12)$$

The logic following (11) leads to the infeasible generalized principal components estimator, $\hat{F}_t(N^{-1}\tilde{\Lambda})$, where $\tilde{\Lambda}$ are the scaled eigenvectors corresponding to the r largest eigenvalues of $\Sigma_e^{-1/2}\hat{\Sigma}_X\Sigma_e^{-1/2}$.

The generalized principal components estimator $\hat{F}_t(N^{-1}\tilde{\Lambda})$ is infeasible because Σ_e is unknown. The challenge in making this estimator feasible is finding a well-behaved estimator of Σ_e . In applications in which N is comparable to or larger than T , the usual estimator of Σ_e based on the residuals is poorly behaved (if $N > T$, it is singular).

At least three versions of feasible generalized principal components estimation have been proposed for the DFM. First, Forni, Hallin, Lippi, and Reichlin (2005) rearrange the decomposition, $\Sigma_X = \Sigma_{\Lambda F} + \Sigma_e$, where $\Sigma_{\Lambda F}$ is the variance of the common component ΛF_t (this decomposition follows from (4)) to obtain $\hat{\Sigma}_e = \hat{\Sigma}_X - \hat{\Sigma}_{\Lambda F}$. They propose estimating $\hat{\Sigma}_{\Lambda F}$ by dynamic principal components (discussed below). Second, Boivin and Ng (2003) suggest a two-step approach using the estimator $\hat{\Sigma}_e = \text{diag}(\{s_{e_t}^2\})$, where $s_{e_t}^2$ is the usual estimator of the error variance of the regression of X_{it} onto the principal components estimator \hat{F}_t ; by setting the off-diagonal terms in Σ_e to zero, their weight matrix has only N estimated elements. Neither of these approaches address possible serial correlation in e_t . To take this into account, Stock and Watson (2005) suggest a three-step approach, akin to the Cochrane-Orcutt estimator, in which F_t is first estimated by principal components, N separate autoregressions are fit to the residuals of the regression of X_{it} on \hat{F}_t , X_{it} is quasi-differenced using the coefficients of the i^{th}

autoregression, and the Boivin-Ng (2003) diagonal Σ_e method is then applied to these quasi-differences.

Dynamic principal components. Dynamic principal components is the frequency domain analog of principal components developed by Brillinger (1964, 1981). Forni, Hallin, Lippi, and Reichlin (2000, 2004) prove the consistency, and provide rates of convergence, of the common component estimated by dynamic principal components. Their method for estimation of f_t by dynamic principal components requires two-sided smoothing, so estimates of f_t at the end of the sample are not available. Consequently dynamic principal components cannot be used directly for forecasting, instrumental variables regression, FAVAR, or other applications that require estimates of f_t for the full sample, and we do not discuss this method further in this survey.

2.3 Third generation: hybrid principal components and state space methods

The third generation of methods for estimating the factors merges the statistical efficiency of the state space approach with the robustness and convenience of the principal components approach. This merger is of particular value for real time applications since the Kalman filter readily handles missing data and can be implemented in real time as individual data are released; for further discussion see Giannone, Reichlin, and Small (2008) and Reichlin's chapter on "Nowcasting" in this Handbook. Additionally, the Kalman filter and smoother average across both series and time, not just across series as in the principal components estimators. As a result, state space/Kalman filter estimates can produce substantial improvements in estimates of the factors and common components if the "signal" of the common component is persistent (so time

averaging helps) and small (so substantial noise remains after cross-series averaging); for an empirical example see Reiss and Watson (2010).

This merged estimation procedure occurs in two steps, which are described in more detail in Giannone, Reichlin, and Small (2008) and Doz, Giannone, and Reichlin (2006). First, the factors are estimated by principal components or generalized principal components. In the second step, these estimated factors \hat{F}_t are used to estimate the unknown parameters of the state space representation. The specifics of how to do this depend on whether the state vector is specified in terms of the static or dynamic factors.

State space model with static factors. In this case, the state space model is given by (4) – (6). Given estimates \hat{F}_t , Λ is estimated by a regression of X_t on \hat{F}_t , and the residuals from this regression are used to estimate the univariate autoregressions in (6). The VAR coefficients $\Psi(L)$ can be estimated by a regression of \hat{F}_t onto its lags, and the variance of $G\eta_t$ can be estimated by the residuals from this VAR.

State space model with dynamic factors. In this case, the state space model is given by (1), (2), and (6), where now f_t and its lags explicitly enter the state vector instead of F_t . Because the dimension q of f_t is less than the dimension r of F_t , the number of parameters is reduced by formulating the state space model in terms of f_t ; that is, the VAR for f_t given in (2) leads to constraints on the lag polynomial $\Phi(L)$ in (5). Given estimates \hat{f}_t of the dynamic factors, computed as described above, the coefficients of (1) can be estimated by regressions of X_t onto \hat{f}_t and its lags; the coefficients of (2) can be estimated by estimating a VAR for \hat{f}_t ; and the coefficients of (6) can be estimated using the residuals from the regression of X_t onto \hat{f}_t .

These estimated parameters fully populate the state space model so that an improved estimate of F_t or f_t , which now invokes time-series averaging, can be computed using the Kalman smoother.

It is also possible to use these estimated coefficients as consistent starting values for maximum likelihood estimation of the coefficients. The MLEs can be computed using the EM algorithm, see for example Engle and Watson (1983), Quah and Sargent (1993) and Doz, Giannone, and Reichlin (2006). Jungbacker and Koopman (2008) show how to speed up the evaluation of the Kalman filter in the DFM by transforming the X_t into an $r \times 1$ vector. Jungbacker, Koopman, and van der Wel (2009) provide additional computational devices that can be used when there are missing data.

2.4 Comparisons of estimators

Several studies have compared the performance of principal components estimators and various feasible generalized principal components estimators in Monte Carlo exercises and in forecast comparisons with actual data. The studies reach somewhat different conclusions when N is small, presumably because the study designs differ. Forni, Hallin, Lippi, and Reichlin (2005) find, in a Monte Carlo study, that their generalized principal components estimator is substantially more precise than the principal components estimator of the common component when there are persistent dynamics in the factors and in the idiosyncratic disturbances, although these differences disappear for large N and T . In contrast, Boivin and Ng (2005), using a Monte Carlo design calibrated to U.S. data, find only minor differences between the principal components and Forni et. al. (2005) generalized principal components estimators.

Boivin and Ng (2005) and D’Agostino and Giannone (2006) also compare the forecasting performance of the principal components estimator to various feasible generalized principal components estimators. Although there are nuances, the overarching conclusion of these comparisons is that the forecasts produced using the various estimators of the factors are highly collinear (holding constant the forecast specification and changing only the factor estimator), and produce very similar R^2 s. There is some evidence that the generalized principal components estimators produce more variable forecasts (sometimes better, sometimes worse) than principal components when N is small, but for values of N and T typical of applied work there is negligible difference in the performance, as measured by pseudo out-of-sample mean squared error, among the forecasts based on the various estimators. This result reassuringly accords with the intuition provided at the beginning of this section, that there is no unique weight matrix W that produces a consistent estimator of the factors, $\hat{F}_t(W)$; rather, sufficient conditions for consistency are that $W'\Lambda \rightarrow H$, where H is full rank, and that $W'\Sigma_e W \rightarrow 0$.

2.5 Bayes estimation

The DFM parameters and factors also can be estimated using Bayesian methods. Historically, there are three main motivations for using Bayes methods to estimate DFMs: first, integration to compute the posterior can be numerically easier and more stable than maximizing the likelihood when there are very many unknown parameters; second, Markov Chain Monte Carlo methods can be used to compute posteriors in nonlinear/nonGaussian latent variable models in which it is exceedingly difficult to compute the likelihood directly; and third, some analysts might wish to impose *a-priori* information on the model in the form of a prior distribution. The first two of these

motivations, however, are less relevant today, at least for DFM applications with large N and relatively few factors: the second-generation nonparametric estimators sidestep the numerical challenges of brute-force MLE and allow the factors and idiosyncratic terms to follow nonlinear and/or nonGaussian processes, and the third-generation methods refine these nonparametric estimates using a high-dimensional parametric state-space model.

Bayes estimation of the DFM parameters and the factor are based on Markov Chain Monte Carlo methods. Chib and Greenberg (1996) lay out the basics of Gibbs sampling applied to linear/Gaussian state space models. Otrok and Whiteman (1996) provide an early implementation of these methods to a linear/Gaussian DFM in state space, in which they estimate a single dynamic factor using four variables. Kose, Otrok, and Whiteman (2003, 2008) use these methods to characterize international comovements and to study international transmission of economic shocks; Kose, Otrok, and Whiteman's (2008) model has 7 common factors among 60 countries, with 3 series per country (so $N = 180$) and $T = 30$ annual observations. Bernanke, Boivin, and Eliasziw (2005) use Gibbs sampling to estimate a state-space version of a Factor Augmented Vector Autoregression (FAVAR, discussed below), but they report that doing so provides only modest changes relative to a second-generation estimation approach in which principle components estimates of the factors are treated as data. Boivin and Giannoni (2006) use Markov Chain Monte Carlo methods to estimate the parameters of a dynamic stochastic general equilibrium (DSGE) model for the latent factors, that is, a model for $\Psi(L)$ and $\text{var}(\eta_t)$ in (2), which are measured by observable series, thus imposing structure on $\lambda(L)$ to identify the factors. In Boivin and Giannoni's (2006) application, and in the

DSGE literature more generally, the priors serve to overcome what appears to be a lack of identification or weak identification of some of the DSGE parameters.

Although the focus of this chapter is the linear DFM, with Gaussian errors when treated parametrically, Bayesian methods are particularly useful when the model contains nonlinear and/or nonGaussian elements. For example, Kim and Nelson (1998) estimate a four-variable DFM in which a single latent factor has a mean that follows a latent Hamilton (1989) Markov switching (regime switching) process. The Hamilton (1989) filter does not extend to this case, that is, for this model, closed-form expressions for the integrals in the general nonlinear/nonGaussian filter in Kitagawa (1987) are not known. The hierarchical nature of this model, however, lends itself to Gibbs sampling and thus to computation of the posterior distribution of the model parameters and the dynamic factor. Carlin, Polson and Stoffer (1992) lay out the general MCMC approach for nonlinear/nonGaussian state space models. It should be noted, however, that in a large- N context, the difficulties posed by the nonlinear/nonGaussian structure for the factors in Kim and Nelson (1998) are also handled by second-generation classical methods. For example, if N is large then \hat{F}_t can be treated as data and the original Hamilton (1989) switching model can be applied instead of using the Kim and Nelson (1998) method, although we are not aware of an application that does so.

A practical motivation for using Bayes methods would be that they produce better forecasts than second- or third-generation classical methods, however we are not aware of a study that undertakes such a comparison.

3. Determining the Number of Factors

Several methods are available for estimating the number of static factors r and the number of dynamic factors q .

3.1 Estimating the number of static factors r

The number of static factors r can be determined by a combination of *a-priori* knowledge, visual inspection of a scree plot, and the use of information criteria developed by Bai and Ng (2002).

Scree plots as a visual diagnostic. A scree plot, introduced by Catell (1966), is a plot of the ordered eigenvalues of $\hat{\Sigma}_X$ against the rank of that eigenvalues. Scree plots are useful diagnostic measures that allow one to assess visually the marginal contribution of the i^{th} principal component to the (trace) R^2 of the regression of X_t against the first i principal components. The scree plot is a useful visual diagnostic; formal tests based on scree plots are discussed below.

Estimation of r based on information criteria. Bai and Ng (2002) developed a family of estimators of r that are motivated by information criteria used in model selection. Information criteria trade off the benefit of including an additional factor (or, more generally, an additional parameter in a model) against the cost of increased sampling variability arising from estimating another parameter. This is done by minimizing a penalized likelihood or log sum of squares, where the penalty factor increases linearly in the number of factors (or parameters). In the DFM case, Bai and Ng (2002) propose minimizing the penalized sum of squares,

$$IC(r) = \ln V_r(\hat{A}, \hat{F}) + rg(N, T), \quad (13)$$

where $V_r(\hat{\Lambda}, \hat{F})$ is the least squares objective function in (11), evaluated at the principal components estimators $(\hat{\Lambda}, \hat{F})$, and where $g(N, T)$ is a penalty factor such that $g(N, T) \rightarrow 0$ and $\min(N, T)g(N, T) \rightarrow \infty$ as $N, T \rightarrow \infty$. These latter conditions are the generalizations to $N, T \rightarrow \infty$ of the standard conditions for the consistency of information criteria in regression, see for example Geweke and Meese (1981). Bai and Ng (2002, 2006b) show that, under the conditions of the approximate DFM, the value of r, \hat{r} , that minimizes an information criterion with $g(N, T)$ satisfying these conditions is consistent for the true value of r , assuming that value of r is finite and does not increase with (N, T) .

A specific choice for $g(N, T)$ that does well in simulations (e.g. Bai and Ng (2002)) is $g(N, T) = (N+T)\ln(\min(N, T))/(NT)$. In the special case $N = T$ this $g(N, T)$ simplifies to two times the familiar BIC penalty factor, that is, $g(T, T) = 2T^{-1}\ln T$. Bai and Ng (2002) refer to the information criterion as IC_{p2} .

Ahn and Horenstein (2009) build on the theoretical results in Bai-Ng (2002) and propose estimating r as the maximizer of the ratio of adjoining eigenvalues; intuitively, this corresponds to finding the edge of the cliff in the scree plot. Their Monte Carlo simulation suggests that this might be a promising new approach that sidesteps a somewhat arbitrary choice about which penalty factor to use in the Bai-Ng (2002) information criterion approach.

Formal tests based on scree plots. Formal distribution theory for the scree plot, and in particular for how to test for whether a seemingly large eigenvalue is in fact large enough to indicate the presence of latent factors, is well known for the special case that, under the null hypothesis, X_{it} is i.i.d. $N(0, 1)$; however a general theory of scree plots has only recently been developed. In the special case that X_{it} is i.i.d. standard normal, then

the eigenvalues are those of a Wishart distribution (see Anderson (1984)). If in addition $N, T \rightarrow \infty$, the largest (centered and rescaled) eigenvalue has the Tracy-Widom (1994) distribution (Johnstone (2001)); this finding means that one need not use the exact Wishart distribution, which depends on (N, T) . El Karoui (2007) generalizes Johnstone (2001) to the case that $\{X_{it}, t = 1, \dots, T\}$ is serially correlated but independent over i , subject to the condition that all series $i = 1, \dots, N$ have the same spectrum which satisfies certain smoothness conditions. Onatski (2008) extends El Karoui (2007) to provide a joint limit (a vector Tracy-Widom law) for the centered and rescaled first several eigenvalues. Because the problem is symmetric in N and T , this result applies equally to panels with cross-sectional correlation but no time series dependence. Onatski (2009) uses the results in Onatski (2008) to develop a pivotal statistic that can be used to test the hypothesis that $q = q_0$ against the hypothesis that $q > q_0$. Onatski's (2009) test is in the spirit of Brillinger's generalized principal components analysis in that it looks at a function of the eigenvalues of the smoothed spectral density matrix at a given frequency, where the function is chosen so that the test statistic is pivotal.

3.2 Estimating the number of dynamic factors q

Three methods have been proposed for formal estimation of the number of dynamic factors. Hallin and Liška (2007) propose a frequency-domain procedure based on the observation that the rank of the spectrum of the common component of X_t is q . Bai and Ng (2007) propose an estimator based on the observation that the innovation variance matrix in the population VAR (5) has rank q . Their procedure entails first estimating the sample VAR by regressing the principal components estimator \hat{F}_t on its lags, then comparing the eigenvalues of the residual variance matrix from this VAR to a

shrinking bound that depends on (N, T) . Amenguel and Watson's (2007) estimator is based on noting that, in a regression of X_t on past values of F_t , the residuals have a factor structure with rank q ; they show that the Bai-Ng (2002) information criterion, applied to the sample variance matrix of these residuals, yields a consistent estimate of the number of dynamic factors. Our own limited Monte Carlo experiments suggest that the Bai-Ng (2007) has somewhat better finite sample performance than the Amenguel-Watson (2007) procedure. We are not aware of a third-party evaluation and comparison of these competing procedures, and such a study is in order.

4. Uses of the Estimated Factors

The estimated factors can be used as data in second-stage regressions and they can be used to estimate structural models, both structural factor-augmented vector autoregressions and DSGEs.

4.1 Use of factors in second stage regressions

Forecasting. As motivated by (3), one step ahead forecasts of a variable y_t (which may or may not be an element of X_t used to estimate the factors) can be computed by regressing y_{t+1} on \hat{F}_t, y_t , lags of y_t , and (optionally) additional lags of \hat{F}_t . Multistep (that is, h -step ahead) forecasts can be computed in two ways. Direct multistep forecasts are computed by regressing y_{t+h} on \hat{F}_t, y_t , and their lags. Alternatively, iterated multistep forecasts can be computed by first estimating a VAR for \hat{F}_t , then using this VAR in conjunction with the one-step ahead forecasting equation to iterate forward h periods. In theory, either direct or iterated forecasts could be better, direct because they avoid potential misspecification error in the VAR for y and F , indirect because they are more

efficient if the VAR model is correctly specified. Empirical evidence provided by Boivin and Ng (2005) for U.S. macro data suggests that the direct method outperforms the iterated method, perhaps (they suggest) because the direct method avoids the risk of misspecification of the process driving the factors. Interestingly, this stands in contrast to Marcellino, Stock, and Watson's (2006) finding that iterated forecasts tend to outperform direct forecasts for models (univariate autoregressions and bivariate ARs) estimated using U.S. macro data. An alternative method for computing iterated forecasts is to iterating forward the state space model, estimated using a hybrid method and the Kalman filter. This has potential advantages over iterating forward a VAR estimated with \hat{F}_t because the dimension of f_t is typically less than F_t which imposes constraints on the VAR that, if correct, reduce sampling error. However we are unaware of systematic empirical evidence comparing iterated forecasts in state space with iterated or direct forecasts based on \hat{F}_t .

Starting with Stock and Watson (1999, 2002b), there is now a very large literature on empirical results of macroeconomic forecasting with high-dimensional DFMs. Eickmeier and Ziegler (2008) conducted a meta-study of 46 distinct forecasting exercises in which DFM forecasts were compared to a variety of benchmarks. The challenges of such a study are considerable, because different studies use different methods, studies vary in quality of execution, the benchmarks differ across studies, and there are other differences. With these caveats, Eickmeier and Ziegler (2008) find mixed results for factor model forecasts, with factor forecasts outperforming competitors in some instances but not others. Some of their findings accord with the econometric theory, for example

the size of the data set positively influences the quality of the factor forecasts, while other findings seem to represent variations across series.

One common finding of pseudo out-of-sample forecasting exercises is that the gains of factor models for forecasting real series tend to be larger than for nominal series, at least for U.S. data. For example, Boivin and Ng (2005) report pseudo out-of-sample relative mean squared forecast errors for factor forecasts, relative to a univariate autoregression benchmark, of 0.55 to 0.83 at the 6-month horizon, and of .49 to .88 at the 12-month horizon, for four major real monthly U.S. macro series (industrial production, employment, real manufacturing and trade sales, and real personal income less transfers), with the range depending on the series and the method for estimating the factors and for computing the multistep forecast. In contrast, they report typical relative mean squared errors of approximately 0.9 for four monthly inflation series at the 6-month horizon, and widely varying results at the 12-month horizon that indicate an undesirable sensitivity of performance to forecast details and to how one measures inflation. The finding that U.S. inflation, and nominal series more generally, can be forecasted by factor models appears to hinge on using data prior to the mid-1980s; it is quite difficult to improve upon simple benchmark models for forecasting U.S. inflation subsequent to the 1990s (e.g. Stock and Watson (2009b)). Eickmeier and Ziegler's (2008) study suggests that the quantitative and even qualitative findings about factor forecast performance in the U.S. do not necessarily generalize to European data.

Stock and Watson (2009c) compare factor forecasts to other high-dimensional forecasts in U.S. data and also find mixed results. For some series, such as real economic activity variables, factor forecasts provide substantial improvements over a range of

small- and large-dimensional competitors, and for these series factor forecasts constitute a success and a significant step forward. But for other series, for example real wage growth, there appears to be valuable forecasting information in principal components beyond the first few used in standard factor model forecasts; for these series, large-model forecasting approaches are valuable compared with small models, but those approaches need to go beyond factor models. There are also some series, such as inflation and exchange rates, which defy all forecasting attempts.

Factors as instrumental variables. Kapetanios and Marcellino (2008) and Bai and Ng (2010) consider the use of estimated factors as instrumental variables. The motivation for so doing is that the factors condense the information in a large number of series, so that one can conduct instrumental variables or generalized method of moments (GMM) analysis using fewer and potentially stronger instruments than if one were to use the original data. The proposal of using the principal components of a large number of series as instruments dates to Klok and Mennes (1960) and Amemiya (1966), however early treatments required strict exogeneity of the instruments. The requirement of strict exogeneity can be weakened if there is a factor structure and if F_t is a valid instrument; for details see Kapetanios and Marcellino (2008) and Bai and Ng (2010).

The main result of Kapetanios and Marcellino (2008) and Bai and Ng (2010) is that, if F_t constitutes a strong instrument and $\hat{\beta}^{TSLS}(F)$ is the two stage least squares estimator based on the instruments F , then under the Bai-Ng (2006a) conditions (in particular, $N^2/T \rightarrow \infty$), $\sqrt{T} \left[\hat{\beta}^{TSLS}(F) - \hat{\beta}^{TSLS}(\hat{F}) \right] \xrightarrow{p} 0$, that is, the principal components estimator can be treated as data for the purpose of instrumental variables regression with

strong instruments. Extensions of this result include the irrelevance of estimation of F even if F_t is a weak instrument, assuming there is a factor structure.

Empirical applications in which estimated factors are used as instruments include Favero, Marcellino, and Neglia (2005) and Beyer, Farmer, Henry, and Marcellino (2008).

4.2 Factor-Augmented Vector Autoregression (FAVAR)

Bernanke, Boivin and Eliasch (2005) introduced the FAVAR as a way to get around two related problems in structural VAR modeling. First, in a conventional (unrestricted) VAR with N variables, the number of parameters per grows with N^2 , so that unrestricted VARs are infeasible when N/T is large. One solution to this problem is to impose structure in the form of a prior distribution on the parameters but that requires formulating a prior distribution. Second, a consequence of using low-dimensional VARs is the possibility that the space of VAR innovations might not span the space of structural shocks, that is, the VAR innovations cannot be inverted to obtain the structural shocks, in which case structural VAR modeling will fail; this failure is called the invertibility or nonfundamentality problem of SVARs.

The key insight of Bernanke, Boivin, and Eliasch (2005) was that the dimensionality problem could be solved by imposing restrictions derived from the DFM. To see this, it is useful to write the static DFM (4) – (6) in VAR form. The result, derived in Stock and Watson (2005), is

$$\begin{pmatrix} F_t \\ X_t \end{pmatrix} = \begin{pmatrix} \Phi(L) & 0 \\ \Lambda\Phi(L) - D(L)\Lambda & D(L) \end{pmatrix} \begin{pmatrix} F_{t-1} \\ X_{t-1} \end{pmatrix} + \begin{pmatrix} G & 0 \\ \Lambda G & I \end{pmatrix} \begin{pmatrix} \eta_t \\ \zeta_t \end{pmatrix} \quad (14)$$

where $D(L) = \text{diag}(d_1(L), \dots, d_N(L))$. Inspection of the coefficient matrices in (14) makes the dimension reduction evident: the number of free parameters is $O(N + r^2)$ but the number of elements of the (unrestricted) VAR matrix is $O((N + r)^2)$. Moreover, all the

parameters of (14) can be estimated by regressions involving X_t , \hat{F}_t , and the residuals from the regression of X_t on \hat{F}_t . In particular, the (population) impulse response function with respect to the innovations η_t is given by $\Gamma(L)$ in $X_t = \Gamma(L)\eta_t + e_t$, where $\Gamma(L) = \Lambda\Phi(L)^{-1}G$ (this is obtained by substituting (5) into (4)). The factor loadings Λ can be estimated by a regression of X_t on \hat{F}_t , the lag polynomial $\Phi(L)$ can be obtained from a VAR estimated using \hat{F}_t , the matrix G can be estimated as the eigenvectors corresponding to the q largest eigenvalues of the $r \times r$ variance matrix of the residuals from the \hat{F}_t VAR (in population this variance matrix has only q nonzero eigenvalues), and η_t can be estimated by $\hat{\eta}_t = \hat{G}'[\hat{F}_t - \hat{\Phi}(L)\hat{F}_{t-1}]$ (for details see Stock and Watson (2005)). This provides sample counterparts to the population innovations η_t , with an arbitrary normalization, and to the impulse response function based on these innovations.

The SVAR modeling exercise now entails imposing sufficient identification restrictions so that the structural shocks (or a single structural shock) can be identified as a linear combination of η_t ; that is, so that a $q \times q$ matrix R (or a row of R) can be identified such that $R\eta_t = \varepsilon_t$, where ε_t are the q structural shocks. Forni, Giannone, Lippi, and Reichlin (2009) show that, under plausible heterogeneity conditions on impulse response functions, such a matrix R will in general exist and will be full rank, that is, the DFM innovations will span the space of structural shocks so a FAVAR will solve the invertibility problem. The matrix R (or individual rows of the matrix) can be identified using extensions of the usual SVAR identification toolkit, including imposing short run restrictions (see for example Bernanke, Boivin, and Elias (2005)); imposing long run restrictions akin to those in Blanchard and Quah (1989) and King, Plosser, Stock, and

Watson (1991); identifying shocks by maximizing the fraction of the variance of common components of one or more variables (Giannone, Reichlin, and Sala (2004)); and imposing sign restrictions (Ahmadi and Uhlig (2009)).

4.3 DSGE estimation using DFMs

Sargent (1989) showed that the DFM can be interpreted as relating multiple indicators to a latent low-dimensional model of the economy. Boivin and Giannoni (2006) extend this concept by associating the dynamic factor evolution equation (2) with a log-linearized DSGE. Accordingly, the elements of f_t correspond to latent economic concepts such as inflation and the output gap. The statistical agencies do not observe these latent concepts, but instead produce multiple measures of them, for example different rates of inflation derived from different price indexes. Similarly, econometricians can produce multiple measures of the unobserved output gap. These observable series constitute X_t , and the elements of f_t are identified by exclusion restrictions in the factor loadings (for example, the multiple observed measures of inflation depend directly on the latent inflation factor but not on the other factors).

By introducing multiple indicators of these latent processes, Boivin and Giannoni (2006) bring additional information to bear on the difficult task of estimation of DSGE parameters. In principle this estimation could be done by MLE, however Boivin and Giannoni (2006) use the Bayesian/Markov Chain Monte Carlo methods common in the DSGE estimation literature.

5. Selected Extensions

This section briefly reviews three extensions of DFM research: DFMs with breaks; DFMs that incorporate cointegration and error correction; and structured DFMs such as hierarchical DFMs.

5.1 Breaks and time-varying parameters

Few papers have considered DFMs with breaks or time-varying parameters. Stock and Watson (2002a) showed that the principal components estimator of the factors is consistent even with certain types of breaks or time variation in the factor loadings. The intuition for this result returns to the idea, introduced in Section 2.1, that the cross-sectional averaging estimator $\hat{F}_t(N^{-1}W)$ is consistent for the space spanned by the factors under relatively weak conditions on W , specifically $N^{-1}W'\Lambda \rightarrow H$; it stands to reason, then, that Λ can break or evolve in some limited fashion and the principal components estimator will remain consistent.

Stock and Watson (2009a) considered the case of a single large break with one set of factors and loadings before the break and another set after. They showed that the full-sample principal components estimator of the factor asymptotically spans the space of the two combined factors. Specifically, the factors for the pre-break subsample will be estimated by one linear combination of the principal components, and the post-break factors will be estimated by another linear combination. The break date need not be known or estimated for this to hold. Accordingly, with a break, the number of full-sample factors can exceed the number of subsample factors both before and after the break. In an empirical application to U.S. macroeconomic data, they find that forecasts

based on full-sample factors can outperform those based on subsample factors, even though a break is found empirically.

Breitung and Eickmeier (2009) propose a test for a single structural break in the factor loadings at an unknown date in a DFM, in the i^{th} equation of the static DFM (4). For example, this can be done by regressing X_{it} on \hat{F}_t and \hat{F}_t interacted with the binary variable that equals 1 for $t > \tau$, (where τ is a hypothesized break date), then computing the Wald statistic testing whether the coefficients on the interacted variables equal zero. Breitung and Eickmeier (2009) find evidence of a change in the factor space in the U.S. in the mid-1980s.

Banerjee, Marcellino, and Masten (2007) provide Monte Carlo results on factor-based forecasting with instability in the factor loadings, and find that large breaks, if undetected, can substantially reduce the performance of full-sample factor-based forecasts. This is consistent with the forecast function changing when the factor loadings change, despite the space being spanned by the factors being consistently estimated. In their setting, it is desirable to estimate the forecasting equation using the post-break sample. They also report an application to data from the EU and from Slovenia, which investigates split-sample instability in the factor forecasts (but not the factor estimates themselves).

5.2 Incorporating cointegration and error correction

All the work discussed until now has assumed that all elements of X_t are integrated of order 0 (I(0)). Typically this involves taking first differences of logarithms for real variables, for example. Elements of X_t can also include other stationary

transformations, including error correction terms; for example it is common for interest rate spreads to be included in X_t .

The econometric theory of principal components estimation of the factors requires modification to cover integrated and cointegrated variables. The basic difficulty can be seen by imagining that the original series are all independent random walks, so that X_t is a vector of standardized independent random walks. These independent walks will be subject to the spurious regression problem (Granger and Newbold (1974)) so without further modification $\hat{\Sigma}_X$ will have a limiting expression (as $T \rightarrow \infty$, N fixed) in terms of a N -dimensional demeaned Brownian motion, and $\hat{\Sigma}_X$ will have many large eigenvalues even though the elements of X_t are independent.

Some recent papers propose techniques designed to estimate and exploit factor structure in the presence of nonstationary factors. Bai (2004) shows that if the factors are $I(1)$ and the idiosyncratic disturbances are $I(0)$ – that is, if the Stock-Watson (1988) common trends representation of cointegrated variables holds – then the space spanned by the factors can be estimated consistently, as can the number of factors. Bai and Ng (2004) provide a procedure for estimating the number of nonstationary factors even if some of the idiosyncratic disturbances are integrated of order one ($I(1)$). Banerjee and Marcellino (2008) use the common trends representation to introduce a factor error correction model. Banerjee, Marcellino, and Masten (2009) provide empirical evidence suggesting that forecasts based on the factor error correction model outperform standard DFM forecasts. A premise of this research is that there is large-scale cointegration in the levels variables or, equivalently, that the spectral density of the transformed variables (X_t as defined for the previous sections) has a rank of only $r \ll N$ at frequency zero.

5.3 Hierarchical DFMs

In certain specialized cases, hierarchical structures arise naturally and can be incorporated into DFMs. Kose, Otrok, and Whiteman (2003) provide an early example of such a model, in which both regional and global factors affect the evolution of output, investment, and consumption in 60 countries. Ng and Moench (2009) provide an application to regional housing prices and Stock and Watson (2010) provide a related application to state-level housing starts. Moench, Ng, and Potter (2009) provide a general formulation of multilevel hierarchical DFMs including extensive discussion of computational issues.

5.4 Outlook

Dynamic factor models have the twin appeals of being grounded in dynamic macroeconomic theory (Sargent (1989), Boivin and Giannoni (2006)) and providing a good first-order description of empirical macroeconomic data, in the sense that a small number of factors explain a large fraction of the variance of many macroeconomic series (Sargent and Sims (1977), Giannone, Reichlin, and Sala (2004), Watson (2004)).

Theoretical econometric research on DFMs over the past decade has made a great deal of progress, and a variety of methods are now available for the estimation of the factors and of the number of factors. Theoretical work is ongoing in many related areas, including weak factor structures, tests for the number of factors, and factor models with instability and breaks.

The bulk of empirical work to date has focused on forecasting. A great deal is now known about the performance of factor forecasts and about best practices for forecasting using factors. Broadly speaking, this research has found that linear factor

forecasts perform well to very well relative to competitors for many, but not all, macroeconomic series. For U.S. real activity series, reductions in pseudo out-of-sample mean squared forecast errors at the two- to four-quarter horizon are often in the range of 20%-40%, although smaller or no improvements are seen for other series, such as U.S. inflation post-1990. Parametric (third-generation) DFMs are also particularly well suited for nowcasting. Work on other empirical applications of DFMs, such as structural VARs and estimation of parameters of DSGEs, is newer and while there are relatively few such applications to date, these constitute promising directions for future empirical research.

References

- Ahmadi, P A., and H. Uhlig (2009), "Measuring the Effects of a Shock to Monetary Policy: A Bayesian Factor-Augmented VAR Approach with Sign Restrictions", manuscript, University of Chicago.
- Ahn, S.C. and A.R. Horenstein (2009), "Eigenvalue Ratio Test for the Number of Factors," manuscript, Arizona State University.
- Amemiya, T. (1966), "On the Use of Principal Components of Independent Variables in Two-Stage Least-Squares Estimation," *International Economic Review* 7, 283-303.
- Amengual, D., and M.W. Watson (2007), "Consistent Estimation of the Number of Dynamic Factors in a Large N and T Panel," *Journal of Business and Economic Statistics*, 91-96.
- Anderson, T.W. (1984), *An Introduction to Multivariate Statistical Analysis, second edition*, New York: Wiley.
- Angelini, E., N. Bańbura, and G. Rünstler (2008), "Estimating and Forecasting the Euro Area Monthly National Accounts from a Dynamic Factor Model," ECB Working Paper No. 593.
- Aruoba, S.B., F.X. Diebold, and C. Scotti (2009), "Real-Time Measurement of Business Conditions," *Journal of Business & Economic Statistics* 27, 417-427.
- Bai, J. (2003), "Inferential Theory for Factor Models of Large Dimensions," *Econometrica*, 71, 135-172.
- Bai, J., and S. Ng (2002), "Determining the Number of Factors in Approximate Factor Models," *Econometrica*, 70, 191-221.

- Bai, J., and S. Ng (2004), "A PANIC Attack on Unit Roots and Cointegration,"
Econometrica, 72, 1127-1177.
- Bai, J., and S. Ng (2006a), "Confidence Intervals for Diffusion Index Forecasts and Inference for Factor-Augmented Regressions," *Econometrica*, 74,1133-1150.
- Bai, J., and S. Ng (2006b), "Determining the Number of Factors in Approximate Factor Models, errata" manuscript, Columbia University.
- Bai, J., and S. Ng (2007), "Determining the Number of Primitive Shocks in Factor Models," *Journal of Business and Economic Statistics* 25, 52-60.
- Bai, J., and S. Ng (2008), "Large Dimensional Factor Analysis," *Foundations and Trends in Econometrics*, 3(2): 89-163.
- Bai, J. and S. Ng (2010), "Instrumental Variable Estimation in a Data-Rich Environment," Forthcoming, *Econometric Theory*
- Banerjee, A., and M. Marcellino (2008) "Factor Augmented Error Correction Models", CEPR WP 6707.
- Banerjee, A., M. Marcellino, and I. Masten (2007), "Forecasting Macroeconomic Variables Using Diffusion Indexes in Short Samples With Structural Change," forthcoming in *Forecasting in the Presence of Structural Breaks and Model Uncertainty*, ed. by D. Rapach and M. Wohar, Elsevier.
- Banerjee, A., M. Marcellino and I. Masten, (2009), Forecasting with Factor-augmented Error Correction Models , mimeo, European University Institute
- Bernanke, B.S., J. Boivin, and P. Elias (2005), "Measuring the Effects of Monetary Policy: A Factor-Augmented Vector Autoregressive (FAVAR) Approach,"
Quarterly Journal of Economics, 120, 387-422.

- Beyer A., Famer R., Henry J. and M. Marcellino (2008) "Factor Analysis in a New-Keynesian Model," *Econometrics Journal*, forthcoming
- Blanchard, O.J., and D. Quah (1989), "Dynamic Effects of Aggregate Demand and Supply Disturbances," *American Economic Review*, 79, 655-673.
- Boivin, J., and M. Giannoni (2006), "DSGE Models in a Data-Rich Environment," NBER Working Paper no. 12772.
- Boivin, J., and S. Ng (2005), "Understanding and Comparing Factor-Based Forecasts," *International Journal of Central Banking* 1, 117-151.
- Boivin J. and S. Ng (2006) "Are More Data Always Better for Factor Analysis?", *Journal of Econometrics*, 132, p. 169-194.
- Brillinger, D.R. (1964), "A Frequency Approach to the Techniques of Principal Components, Factor Analysis and Canonical Variates In the Case of Stationary Time Series," Invited Paper, Royal Statistical Society Conference, Cardiff Wales. (Available at <http://stat-www.berkeley.edu/users/brill/papers.html>).
- Brillinger, D.R. (1981), *Time Series: Data Analysis and Theory, expanded edition*, San Francisco: Holden-Day.
- Breitung, J. and S. Eickmeier (2009), "Testing for Structural Breaks in Dynamic Factor Models," Deutsche Bundesbank Economic Studies Discussion Paper No. 05/2009.
- Carlin, B.P., N.G. Polson, and D.S. Stoffer (1992), "A Monte Carlo Approach to Nonnormal and Nonlinear State-Space Modeling," *Journal of the American Statistical Association*, 87, 493-500.

- Cattell, R. B. (1966) "The Scree Test for the Number of Factors", *Multivariate Behavioral Research*, vol. 1, 245-76.
- Chamberlain, G., and M. Rothschild (1983), "Arbitrage Factor Structure, and Mean-Variance Analysis of Large Asset Markets," *Econometrica*, 51,1281-1304.
- Chib, S. and E. Greenberg (1996), "Markov Chain Monte Carlo Simulation Methods in Econometrics," *Econometric Theory*, 12, 409-431.
- Connor, G., and R.A. Korajczyk (1986), "Performance Measurement with the Arbitrage Pricing Theory," *Journal of Financial Economics*, 15, 373-394.
- D'Agostino, A., and Domenico Giannone (2006), "Comparing Alternative Predictors Based on Large-Panel Factor Models," ECB Working Paper 680.
- Doz, C., D. Giannone, and L. Reichlin (2006), "A Quasi Maximum Likelihood Approach for Large Approximate Dynamic Factor Models," ECB Working Paper 674.
- Eickmeier, S., C. Ziegler (2008), How successful are dynamic factor models at forecasting output and inflation? A meta-analytic approach, *Journal of Forecasting*, 27(3), 237-265.
- El Karoui, N. (2007), "Tracy-Widom Limit for the Largest Eigenvalue of a Large Class of Complex Sample Covariance Matrices," *Annals of Probability*, 35, 633-714.
- Engle, R.F., and M.W. Watson (1981), "A One-Factor Multivariate Time Series Model of Metropolitan Wage Rates," *Journal of the American Statistical Association*, 76, 774-781.
- Engle, R.F., and M.W. Watson (1983), "Alternative Algorithms for Estimation of Dynamic MIMIC, Factor, and Time Varying Coefficient Regression Models," *Journal of Econometrics*, Vol. 23, pp. 385-400.

- Favero, C.A., M. Marcellino, and F. Neglia (2005), “Principal Components at Work: The Empirical Analysis of Monetary Policy With Large Datasets,” forthcoming, *Journal of Applied Econometrics*.
- Forni, M., D. Giannone, M. Lippi, and L. Reichlin (2009), “Opening the Black Box: Structural Factor Models with Large Cross Sections,” *Econometric Theory*, 25, 1319-1347.
- Forni, M., M. Hallin, M. Lippi, and L. Reichlin (2000), “The Generalized Factor Model: Identification And Estimation,” *Review of Economics and Statistics*, 82, 540–554.
- Forni, M., M. Hallin, M. Lippi, and L. Reichlin (2004), “The Generalized Factor Model: Consistency and Rates,” *Journal of Econometrics*, 119, 231-255.
- Forni, M., M. Hallin, M. Lippi, and L. Reichlin (2005), “The Generalized Dynamic Factor Model: One-Sided Estimation and Forecasting,” *Journal of the American Statistical Association*, 100, 830-839.
- Forni, M., and L. Reichlin (1998), “Let’s Get Real: A Dynamic Factor Analytical Approach to Disaggregated Business Cycle,” *Review of Economic Studies*, 65, 453-474.
- Geweke, J. (1977), “The Dynamic Factor Analysis of Economic Time Series,” in *Latent Variables in Socio-Economic Models*, ed. by D.J. Aigner and A.S. Goldberger, Amsterdam: North-Holland.
- Geweke, J. and R. Meese (1981), “Estimating Regression Models of Finite but Unknown Order,” *International Economic Review* 22, no. 1, 55-70.
- Giannone, D., L. Reichlin, and L. Sala (2004), “Monetary Policy in Real Time,” *NBER Macroeconomics Annual*, 2004, 161-200.

- Giannone, D., L. Reichlin, and D. Small (2008), “Nowcasting: The Real-Time Informational Content of Macroeconomic Data,” *Journal of Monetary Economics*, 55, 665-676.
- Granger, P. and C.W.J. Newbold (1974), “Spurious Regressions in Econometrics,” *Journal of Econometrics* 2, 111-120.
- Hallin, M., and Liška, R., (2007), “The Generalized Dynamic Factor Model: Determining the Number of Factors,” *Journal of the American Statistical Association*, 102, 603-617.
- Hamilton, J.D., (1989), “A New Approach to the Economic Analysis of Nonstationary Time Series and the Business Cycle,” *Econometrica*, 57, 357-384.
- Harvey, A.C. (1989), *Forecasting, Structural Time Series Models and the Kalman Filter*. Cambridge, U.K.: Cambridge University Press, 1989
- Johnstone, I.M. (2001), “On the Distribution of the Largest Eigenvalue in Principal Component Analysis,” *Annals of Statistics*, 29, 295-327.
- Jungbacker, D., and S.J. Koopman (2008), “Likelihood-Based Analysis for Dynamic Factor Models,” manuscript, Tinbergen Institute.
- Jungbacker, D., S.J. Koopman, and M. van der Wel (2009), “Dynamic Factor Analysis in the Presence of Missing Data,” Tinbergen Institute Discussion Paper 2009-010/4.
- Kapetanios, G., and M. Marcellino (2008), “Factor-GMM Estimation with Large Sets of Possibly Weak Instruments,” manuscript, EUI
- Kim, C-J., and C. Nelson (1998), “Business Cycle Turning Points, A New Coincident Index, and Tests of Duration Dependence Based on a Dynamic Factor Model with Regime-Switching,” *Review of Economics and Statistics*, 80(2), 188-201.

- King, Robert G., C.I. Plosser, J.H. Stock, and M.W. Watson (1991), "Stochastic Trends and Economic Fluctuations," *American Economic Review* 81(4): 819-840.
- Kitagawa, G., (1987), "Non-Gaussian State Space Modeling of Non-Stationary Time Series," with discussion, *Journal of the American Statistical Association*, 82, 1032-1063.
- Kloek, T., and L.B.M. Mennes (1960), "Simultaneous Equations Estimation Based on Principal Components of Predetermined Variables," *Econometrica*, 28, 45-61.
- Kose, A., C. Otrok, C., and C. Whiteman, (2003), "International Business Cycles: World Region and Country Specific Factors," *American Economic Review* 93:4, 1216-1239.
- Kose, A., C. Otrok, C., and C. Whiteman, (2008), "Understanding the Evolution of World Business Cycles," *Journal of International Economics*, 75, 110-130.
- Marcellino, M., J.H. Stock, and M.W. Watson, (2006), "A Comparison of Direct and Iterated Multistep AR Methods for Forecasting Macroeconomic Series," *Journal of Econometrics*, 135, 499-526.
- Moench E. and S. Ng and S. Potter (2009), "Dynamic Hierarchical Factor Models," manuscript, Columbia University.
- Ng, S. and E. Moench (2009), "A Factor Analysis of Housing Market Dynamics in the U.S. and the Regions," manuscript, Columbia University.
- Onatski, A. (2008), "The Tracy-Widom Limit for the Largest Eigenvalues of Singular Complex Wishart Matrices," *Annals of Applied Probability* 18, 470-490.
- Onatski, Alexei (2009), "Testing Hypotheses about the Number of Factors in Large Factor Models," *Econometrica* 77 1447-1479.

- Otrok, C., and C.H. Whiteman, (1998), "Bayesian Leading Indicators: Measuring and Predicting Economic Conditions in Iowa," *International Economic Review*, 39(4).
- Quah, D., and T.J. Sargent (1993), "A Dynamic Index Model for Large Cross Sections" (with discussion), in *Business Cycles, Indicators, and Forecasting*, ed. by J.H. Stock and M.W. Watson, Chicago: University of Chicago Press for the NBER, 285-310.
- Reiss, R., and M.W. Watson (2010), "Relative Goods' Prices and Pure Inflation," *American Economic Journal: Macroeconomics*, forthcoming.
- Sargent, T.J. (1989), "Two Models of Measurements and the Investment Accelerator," *Journal of Political Economy* 97:251–287.
- Sargent, T.J., and C.A. Sims (1977), "Business Cycle Modeling Without Pretending to Have Too Much A-Priori Economic Theory," in *New Methods in Business Cycle Research*, ed. by C. Sims et al., Minneapolis: Federal Reserve Bank of Minneapolis.
- Stock, J.H. and M.W. Watson (1988), "Testing for Common Trends," *Journal of the American Statistical Association* 83, no. 404, 1097-1107.
- Stock, J.H., and M.W. Watson (1989), "New Indexes of Coincident and Leading Economic Indicators," *NBER Macroeconomics Annual 1989*, 351-393.
- Stock, J.H., and M.W. Watson (1999), "Forecasting Inflation," *Journal of Monetary Economics*, 44, 293-335.
- Stock, J.H., and M.W. Watson (2002a), "Forecasting Using Principal Components from a Large Number of Predictors," *Journal of the American Statistical Association*, 97, 1167-1179.

- Stock, J.H., and M.W. Watson (2002b), "Macroeconomic Forecasting Using Diffusion Indexes," *Journal of Business and Economic Statistics*, 20, 147-162.
- Stock, J.H., and M.W. Watson (2005), "Implications of Dynamic Factor Models for VAR Analysis," manuscript.
- Stock, J.H., and M.W. Watson (2006), "Forecasting with Many Predictors," ch. 6 in *Handbook of Economic Forecasting*, ed. by Graham Elliott, Clive W.J. Granger, and Allan Timmermann, Elsevier, 515-554.
- Stock, J.H., and M.W. Watson (2009a), "Forecasting in Dynamic Factor Models Subject to Structural Instability," Ch. 7. in Neil Shephard and Jennifer Castle (eds), *The Methodology and Practice of Econometrics: Festschrift in Honor of D.F. Hendry*. Oxford; Oxford University Press.
- Stock, J.H., and M.W. Watson (2009b), "Phillips Curve Inflation Forecasts," Ch. 3 in *Understanding Inflation and the Implications for Monetary Policy* (2009), Jeffrey Fuhrer, Yolanda Kodrzycki, Jane Little, and Giovanni Olivei (eds). Cambridge: MIT Press.
- Stock, J.H., and M.W. Watson (2009c), "Generalized Shrinkage Methods for Forecasting Using Many Predictors," manuscript.
- Stock, J.H., and M.W. Watson (2010), "The Evolution of National and Regional Factors in U.S. Housing Construction," Ch. 3 in T. Bollerslev, J. Russell, and M. Watson (eds), *Volatility and Time Series Econometrics: Essays in Honor of Robert F. Engle*, Oxford; Oxford University Press.
- Tracy, C.A., and H. Widom (1994), "Level Spacing Distributions and the Airy Kernel," *Communications in Mathematical Physics*, 159, 151-174.

Watson, M.W. (2004), "Comment on Giannone, Reichlin, and Sala," *NBER Macroeconomics Annual*, 2004, 216-221.