

## OBSERVING IMBH–IMBH BINARY COALESCENCES VIA GRAVITATIONAL RADIATION

JOHN M. FREGEAU<sup>1</sup>, SHANE L. LARSON<sup>2</sup>, M. COLEMAN MILLER<sup>3,4</sup>, RICHARD O’SHAUGHNESSY<sup>1</sup>, FREDERIC A. RASIO<sup>1</sup>

<sup>1</sup>Department of Physics and Astronomy, Northwestern University, Evanston, IL 60208; fregeau@alum.mit.edu, oshaughn@northwestern.edu, rasio@northwestern.edu

<sup>2</sup>Center for Gravitational Wave Physics, The Pennsylvania State University, University Park, PA 16802; shane@gravity.psu.edu

<sup>3</sup>University of Maryland, Department of Astronomy, College Park, MD 20742; miller@astro.umd.edu and

<sup>4</sup>Goddard Space Flight Center, Greenbelt, MD

*Accepted for publication in ApJ*

### ABSTRACT

Recent numerical simulations have suggested the possibility of forming double intermediate mass black holes (IMBHs) via the collisional runaway scenario in young dense star clusters. The two IMBHs formed would exchange into a common binary shortly after their birth, and quickly inspiral and merge. Since space-borne gravitational wave (GW) observatories such as LISA will be able to see the late phases of their inspiral out to several Gpc, and LIGO will be able to see the merger and ringdown out to similar distances, they represent potentially significant GW sources. In this Letter we estimate the rate at which LISA and LIGO will see their inspiral and merger in young star clusters, and discuss the information that can be extracted from the observations. We find that LISA will likely see tens of IMBH–IMBH inspirals per year, while advanced LIGO could see  $\sim 10$  merger and ringdown events per year, with both rates strongly dependent on the distribution of cluster masses and densities.

*Subject headings:* stellar dynamics — black hole physics — gravitational waves

### 1. INTRODUCTION

Observations suggesting the existence of intermediate-mass black holes (IMBHs) have mounted in recent years. Ultra-luminous X-ray sources (ULXs)—point X-ray sources with inferred luminosities  $\gtrsim 10^{39}$  erg/s—may be explained by sub-Eddington accretion onto BHs more massive than the maximum of  $\sim 10M_{\odot}$  expected from stellar core collapse (Miller & Colbert 2004). Similarly, the cuspy velocity dispersion profiles in the centers of the globular clusters M15 and G1 may also be explained by the dynamical influence of a central IMBH (van der Marel et al. 2002; Gerssen et al. 2002; Gebhardt et al. 2005), although this conclusion remains somewhat controversial (Baumgardt et al. 2003).

The most likely formation scenario for an IMBH is the collapse of a very massive star (VMS), which was formed early in the lifetime of a young star cluster via a runaway sequence of physical collisions of massive main-sequence stars (Portegies Zwart et al. 1999; Ebisuzaki et al. 2001; Portegies Zwart & McMillan 2002; Gürkan et al. 2004). This scenario has been studied in detail for star clusters without primordial binaries, with recent work showing that runaway growth of a VMS to  $\sim 10^3 M_{\odot}$  occurs generically in clusters with deep core collapse times shorter than the  $\sim 3$  Myr main-sequence lifetime of the most massive stars (Freitag et al. 2006).

Due to the computational cost of simulating the more realistic case of star clusters with primordial binaries, it is only recently that such simulations have been performed (Portegies Zwart et al. 2004; Gürkan et al. 2006). The work of Gürkan et al. (2006) was the first to systematically study the influence of primordial binaries on the runaway growth process. They showed that stellar collisions during binary scattering interactions offer an alternate channel for runaway growth, with the main result that clusters with binary fractions larger than  $\approx 10\%$  generically produce *two* VMSs, provided the cluster is

sufficiently dense and/or centrally concentrated to trigger the runaway earlier than  $\sim 3$  Myr in the absence of primordial binaries. Observations and recent numerical calculations suggest that star clusters may be born with large binary fractions ( $\gtrsim 30\%$ ; Hut et al. 1992; Ivanova et al. 2005), implying that *all* sufficiently dense and massive star clusters could form multiple VMSs.

The VMSs formed will undergo core-collapse supernovae and likely become IMBHs on a timescale of  $\sim 4$  Myr after cluster formation (the lifetime of a VMS is extended slightly by collisional rejuvenation; see, e.g. Freitag et al. 2006). After their separate formation, the two IMBHs will quickly exchange into a common binary via dynamical interactions. The IMBH–IMBH binary (IMBHB) will then shrink via dynamical friction due to the cluster stars, on a timescale  $\sim t_r \langle m \rangle / M_{\text{IMBH}} \lesssim 10$  Myr, where  $t_r$  is the core relaxation time,  $\langle m \rangle$  is the local average stellar mass, and  $\langle m \rangle / M_{\text{IMBH}} \lesssim 10^{-2}$ . Note that since  $t_r$  scales inversely with  $\langle m \rangle$  for fixed core velocity dispersion and mass density, the dynamical friction timescale is independent of  $\langle m \rangle$  (Binney & Tremaine 1987). The IMBHB will then shrink further via dynamical encounters with cluster stars (Quinlan 1996; Yu & Tremaine 2003; Miller 2005), until it merges quickly via gravitational radiation, on a timescale  $\approx 1 \text{ Myr } (\sigma_c / 20 \text{ km s}^{-1})^3 (\rho_c / 10^5 M_{\odot} \text{ pc}^{-3})^{-1} (M_{\text{IMBH}} / 10^3 M_{\odot})^{-1}$ , where  $\sigma_c$  is the cluster core velocity dispersion and  $\rho_c$  is the core mass density (Quinlan 1996, eqs. [29] and [30]). This timescale has also been confirmed by numerical scattering calculations (Gültekin, private communication).

Only the more massive IMBHBs merge in the LISA band of  $10^{-4}$ –1 Hz (redshifted binary mass  $M_z \equiv (1+z)M \gtrsim 4 \times 10^3 M_{\odot}$ , where  $M$  is the total binary mass). Fig. 1 shows the final gravitational wave (GW) frequency  $f_f$  (the frequency at the inner-most stable circular orbit if within the LISA frequency range (large  $M_z$ ), otherwise




FIG. 1.— The final GW frequency  $f_f$  (see text), and the frequency 1 yr prior,  $f_i$ , for an IMBHB with total mass  $M$  and reduced mass parameter  $\eta$ , as a function of redshifted binary mass  $M_z$ , for  $\eta = 0.25$  (equal-mass binary) and  $\eta = 0.1$  (mass ratio 0.13). (The final frequency is roughly independent of  $\eta$ .)




FIG. 2.— Luminosity distance,  $d_L(z)$ , to which an IMBHB of total mass  $M$  and reduced mass parameter  $\eta$  can be seen via its inspiral with LISA with  $S/N = 10$  for a 1 yr integration, and via its merger and ringdown with  $S/N = 8$  for iLIGO and AdLIGO, as a function of the redshifted mass  $M_z$ . The corresponding redshift (calculated using the WMAP year 3 cosmological parameters, as discussed in the text) is shown on the right vertical axis.

the maximum LISA frequency of  $\approx 1$  Hz (small  $M_z$ , as in Will (2004)), and the frequency 1 yr prior,  $f_i$ , as a function of redshifted mass  $M_z$ , for the reduced mass parameters  $\eta = 0.25$  (equal-mass binary) and  $\eta = 0.1$  (mass ratio 0.13) (see, e.g., Will 2004). For a wide range in  $M_z$ , the late stages of inspiral clearly span the LISA “sweet spot” (roughly a decade centered on  $10^{-2.2}$  Hz), implying that LISA could easily detect the chirp signal, enabling a measurement of the masses of the binary members. Such an observation would be direct evidence for an IMBH.

In Sec. 2 we estimate the rate at which LISA will observe inspiral of IMBHBs in young star clusters. In Sec. 3 we estimate the rate at which LIGO will observe their merger and ringdown. Finally, in Sec. 4 we discuss the observational consequences.

## 2. ESTIMATING THE LISA DETECTION RATE

We first need to know the distance to which LISA can see IMBHB inspirals. Following the techniques in Will (2004) and Flanagan & Hughes (1998), we adopt the latest LISA sensitivity curve (Larson 2003), including confusion noise from Galactic white dwarf binaries (Bender & Hils 1997), and calculate the maximum luminosity distance,  $d_L(z)$ , to which an IMBHB of total mass  $M$  and reduced mass parameter  $\eta$  can be seen with  $S/N = 10$  for a 1 yr integration. The results are shown in Fig. 2 as a function of  $M_z$ , for  $\eta = 0.25$  and  $\eta = 0.1$ . Note that the results of Gürkan et al. (2006) show that the masses of the IMBHBs never differ by more than a factor of a few ( $\eta \gtrsim 0.15$ ). Thus LISA will be able to see typical IMBHBs ( $M \sim 10^3 M_\odot$ ) out to a few Gpc.

With this information in hand, we first make a crude estimate of the LISA event rate. Following Miller (2002), we write for the total rate

$$R \equiv \frac{dN_{\text{event}}}{dt} = \left( \int_0^{z_{\text{max}}} \frac{dV_c}{dz} dz \right) \frac{dN_{\text{cl}}}{dV} g \frac{1}{t_U}. \quad (1)$$

The first factor,  $\int_0^{z_{\text{max}}} (dV_c/dz) dz$ , is the integrated comoving volume of space in which LISA is sensitive to the events. The second factor,  $dN_{\text{cl}}/dV$ , is the number density of star clusters sufficiently massive to form IMBHBs. Since the *globular* clusters we currently see were likely at least a few times more massive at formation (Joshi et al. 2001), we set this factor to the current density of globular clusters in the local universe,  $dN_{\text{cl}}/dV \approx 8h^3 \text{Mpc}^{-3}$  (Portegies Zwart & McMillan 2000). The third factor,  $g$ , is the fraction of sufficiently massive clusters that have a large enough binary fraction and initial central density to produce IMBHBs. Since initial cluster structural parameters are largely unknown, we treat  $g$  as a parameter. The fourth factor is the event rate per IMBHB-producing cluster, taken to be one divided by the age of the universe, since only one IMBHB is formed per cluster over its lifetime. We adopt a  $\Lambda$ CDM cosmology, with parameters  $\Omega_M = 0.24$ ,  $\Omega_\Lambda = 0.76$ , and  $h = 0.73$ , for which  $t_U = 13.8$  Gyr (Spergel et al. 2006). Putting this together for  $d_L = 4.9$  Gpc ( $z_{\text{max}} = 0.79$ ), the distance to which LISA can see IMBHBs with  $M = 2 \times 10^3 M_\odot$ , eq. (1) gives  $R \approx 1(g/0.1) \text{yr}^{-1}$ .

Writing down a generalized form of the rate integral in eq. (1) is straight-forward. Since the time between cluster formation and IMBHB merger is  $\ll t_U$ , we assume that the merger is coincident with cluster formation. Thus the rate integral is

$$R \equiv \frac{dN_{\text{event}}}{dt_o} = \int_0^{z_{\text{max}}} \frac{d^2 M_{\text{SF}}}{dV_c dt_e} g_{\text{cl}} g \times \frac{dt_e}{dt_o} \frac{dV_c}{dz} \int_{M_{\text{cl}, \text{min}}(z)}^{M_{\text{cl}, \text{max}}} \frac{d^2 N_{\text{cl}}}{dM_{\text{SF}, \text{cl}} dM_{\text{cl}}} dM_{\text{cl}} dz. \quad (2)$$

Here  $R \equiv dN_{\text{event}}/dt_o$  is the event rate observed at  $z = 0$  by LISA,  $d^2 M_{\text{SF}}/dV_c dt_e$  is the star formation rate (SFR) in mass per unit of comoving volume per unit of local time,  $g_{\text{cl}}$  is the fraction of star forming mass that goes into star clusters more massive than  $10^{3.5} M_\odot$  (generally a function of  $z$ ),  $g$  is as above, and  $d^2 N_{\text{cl}}/dM_{\text{SF}, \text{cl}} dM_{\text{cl}}$  is the distribution function of clusters over individual cluster mass  $M_{\text{cl}}$  and total star forming mass in clusters  $M_{\text{SF}, \text{cl}}$ . Finally,  $dt_e/dt_o$  is simply  $(1+z)^{-1}$ , and  $dV_c/dz$  is the rate of change of comoving volume with

redshift, which is a function of cosmological parameters (Hogg 1999). Note that we set  $z_{\max} = 5$ , since this is roughly the limit to which the cosmic SFR can be traced. Thus the integral in eq. (2) should be considered a mild lower limit to the true rate. We now discuss each element in eq. (2) in more detail.

Following Porciani & Madau (2001), we adopt three different choices for the SFR:

$$\left(\frac{d^2 M}{dV_c dt}\right)_{\text{SF}i} = C_i h_{65} F(z) G_i(z) M_{\odot} \text{yr}^{-1} \text{Mpc}^{-3}, \quad (3)$$

with  $i = 1, 2, 3$  denoting the different rates,  $C_i$  a constant,  $G_i(z)$  a function of  $z$ ,  $h_{65} = h/0.65$ , and  $F(z) = [\Omega_M(1+z)^3 + \Omega_k(1+z)^2 + \Omega_{\Lambda}]^{1/2}/(1+z)^{3/2}$ . The first is from Madau & Pozzetti (2000), with  $C_1 = 0.3$  and  $G_1(z) = e^{3.4z}/(e^{3.8z} + 45)$ , which peaks between  $z = 1$  and 2 and decreases at larger redshift. The second is from Steidel et al. (1999), with  $C_2 = 0.15$  and  $G_2(z) = e^{3.4z}/(e^{3.4z} + 22)$ , which is roughly constant for  $z \gtrsim 2$ . The third is from Blain et al. (1999), with  $C_3 = 0.2$  and  $G_3(z) = e^{3.05z-0.4}/(e^{2.93z} + 15)$ , which increases above  $z \approx 2$ .

Measuring the fraction of star-forming mass in clusters is difficult for anywhere but the local universe. Similarly, while we know reasonably well the initial cluster conditions required to form an IMBHB (Gürkan et al. 2006), we know much less well the distribution of cluster properties at birth. We therefore treat  $g_{\text{cl}}$  and  $g$  as parameters, taking  $g_{\text{cl}} = 0.1$  and  $g = 0.1$  somewhat arbitrarily as canonical values.

Assuming that the spectrum of cluster masses is neither a function of cosmic epoch nor the total star forming mass available for clusters, the factor  $d^2 N_{\text{cl}}/dM_{\text{SF},\text{cl}} dM_{\text{cl}}$  can be separated as

$$\frac{d^2 N_{\text{cl}}}{dM_{\text{SF},\text{cl}} dM_{\text{cl}}} = \frac{f(M_{\text{cl}})}{\int M_{\text{cl}} f(M_{\text{cl}}) dM_{\text{cl}}}, \quad (4)$$

where  $f(M_{\text{cl}})$  is the (normalized) distribution function of cluster masses. For this we adopt the power-law form observed for young star clusters in the Antennae, which is thought to be universal:  $f(M_{\text{cl}}) \propto M_{\text{cl}}^{-2}$  (Zhang & Fall 1999), with a lower limit of  $10^{3.5} M_{\odot}$  and an upper limit of  $10^7 M_{\odot}$ .

It is the limit  $M_{\text{cl},\min}(z)$  in eq. (2) that encodes all information about the detectability of an IMBHB inspiral by LISA. Specifically, the redshift to which LISA can see the inspiral is a function of the binary mass, which is itself a function of the host cluster mass. Adopting an efficiency factor  $f_{\text{GC}}$  for the fraction of cluster mass going into the IMBHB, this relationship is inverted to obtain  $M_{\text{cl},\min}(z)$ . Recent numerical work shows that the efficiency factor is  $f_{\text{GC}} \approx 2 \times 10^{-3}$ , independent of cluster initial conditions (Gürkan et al. 2004), which we take as our canonical value. At low redshift,  $M_{\text{cl},\min}(z)$  is clamped at the value  $M_{\text{cl}} = 200 M_{\odot}/f_{\text{GC}}$ , set by adopting the definition that an IMBH have mass  $\geq 10^2 M_{\odot}$ . At high redshift ( $z > 5$ , so not relevant to our calculation),  $M_{\text{cl},\min}(z)$  is clamped at the value of  $10^7 M_{\odot}$  from the cluster mass function; in other words no cluster is sufficiently massive to produce an IMBHB massive enough to be observable by LISA, so the integral is zero.

We numerically integrated eq. (2) for the different SFRs in eqs. (3), for  $S/N = 10$  and an integration time




FIG. 3.— Integrand of the rate integral in eq. (2) for the three different SFRs in eqs. (3), for  $\eta = 0.25$  and  $\eta = 0.1$ .

of 1 yr, to find that the rate is

$$R(\eta = 0.25) \approx 40\text{--}50 \left(\frac{g_{\text{cl}}}{0.1}\right) \left(\frac{g}{0.1}\right) \text{yr}^{-1}, \quad (5)$$

with the spread in the coefficient from the different SFRs. The coefficient decreases to 20–25 for  $\eta = 0.1$ . The rate is dominated by clusters in the mass range  $10^6\text{--}10^{6.5} M_{\odot}$  (IMBHB mass  $2 \times 10^3\text{--}6 \times 10^3 M_{\odot}$ ), with more than half the contribution to the rate coming from this mass range, for both  $\eta = 0.1$  and  $\eta = 0.25$ , and for all three SFRs in eq. (3) (except SF3 for  $\eta = 0.1$ ). Note that eq. (2) is only strictly valid when the source is visible by the instrument for less than the integration time. This turns out not to be precisely correct. A typical IMBHB with mass  $M = f_{\text{GC}} 10^{6.25} M_{\odot}$  takes roughly 4 years to cross the LISA band from the edge of the white dwarf confusion knee at  $\approx 2$  mHz to the upper edge of the band at  $\approx 1$  Hz. Thus the rate presented in eq. (5) is an underestimate by of order a factor of a few.

Fig. 3 shows the integrand of the rate integral in eq. (2) for the three different SFRs in eqs. (3), for  $\eta = 0.25$  and  $\eta = 0.1$ . Most events originate from  $z \sim 1$ . Unfortunately, neither  $R$  nor  $dR/dz$  is particularly sensitive to the cosmic SFR, with  $dR/dz$  decreasing quickly above  $z \approx 2$  even when the SFR is increasing (as in SF3). Thus observations of IMBHB inspirals will not be very informative about the cosmic SFR. However, they will likely yield a handle on the fraction of star formation that is in compact massive clusters.

### 3. ESTIMATING THE LIGO DETECTION RATE

Shortly after the two IMBHs merge, the merger product can be well described as a single perturbed black hole, emitting GWs at its quasinormal frequencies. Largely falling within the initial and advanced LIGO (iLIGO and AdLIGO) sensitivity bands, the merger and ring-down waves will likely carry a few percent of the rest mass energy of the hole (see, e.g. Flanagan & Hughes 1998). Numerical simulations suggest that a merging pair of nonspinning equal-mass black holes will emit a fraction  $\epsilon \simeq 0.03$  of their rest mass in merger and ring-down GWs, forming a black hole with spin parameter  $a \simeq 0.7$  (Baker et al. 2002; Campanelli et al. 2006; Baker

et al. 2006). Under these conditions, the ringdown frequency is given by (see eq. [3.17] of Flanagan & Hughes 1998)

$$f \approx \frac{c^3}{2\pi GM_z} (1 - 0.63(1-a)^{3/10}) \approx 180 \left( \frac{M_z}{10^2 M_\odot} \right)^{-1} \text{ Hz.} \quad (6)$$

We can express the distance to which we are sensitive to ringdown waves at signal-to-noise ratio  $\rho$  as

$$d_L(z) = \left( \frac{2\epsilon M_z}{5\pi^2 \rho^2 f^2 S(f)} \right)^{1/2}, \quad (7)$$

where  $S(f)$  is the spectral noise density of LIGO. Combining this expression with the concordance cosmological model and iLIGO and AdLIGO sensitivity curves, we find the range to which LIGO can detect ringdown shown in Fig. 2.

To obtain a conservative estimate for the rate at which iLIGO and AdLIGO could detect these mergers with a ringdown-only search, we use eq. (1) with a moderately optimistic range of  $d_L \approx 100$  Mpc for iLIGO and  $d_L = 2$  Gpc for AdLIGO. The expected detection rate is then  $10^{-4}(g/0.1) \text{ yr}^{-1}$  and  $1(g/0.1) \text{ yr}^{-1}$ , for iLIGO and AdLIGO, respectively. More detailed estimates using machinery analogous to eq. (2) increase these estimates by roughly an order of magnitude, making the rate for AdLIGO  $10(g_{\text{cl}}/0.1)(g/0.1) \text{ yr}^{-1}$ .

#### 4. DISCUSSION

It appears likely that LISA will see tens of IMBHB inspiral events per year, while AdLIGO could see  $\sim 10$  merger and ringdown events per year, with both rates strongly dependent on the distribution of cluster masses and densities. Detection of an IMBHB would have profound implications. A match-filtered observation of the

inspiral would yield the redshifted masses of the black holes, directly confirming the existence of IMBHBs. It would also yield the luminosity distance to the source; with enough observations, constraints could be placed on the cosmic history of star formation in dense, massive clusters. Detection of the ringdown signal from the merger product will additionally yield its spin, which may provide insight into its formation history.

Typical IMBHBs spend  $\gtrsim 10^6$  yr inspiraling through the LISA band, with nearly all of that time spent at low frequencies ( $\lesssim 10^{-3}$  Hz). In the low frequency region they will thus appear as a large number of monochromatic sources, possibly contributing to confusion noise and increasing the noise floor (e.g., Farmer & Phinney 2003). A detailed calculation of this is beyond the scope of this Letter. However, we note that if their contribution is similar in magnitude to that of Galactic compact object binaries (Bender & Hils 1997), the rates predicted in eq. (5) would decrease by about 20%.

We thank Kayhan Gültekin for providing the numerically-calculated timescale for an IMBHB to shrink by stellar dynamical encounters after formation. We also thank the anonymous referee for many helpful comments on the manuscript. JMF and FAR acknowledge support from NASA grant NNG04G176G and NSF grant PHY-0245028. SLL acknowledges support from NASA award NNG05GF71G. MCM was supported in part by NASA grant NAG 5-13229 and by the Research Associateships Programs Office of the National Research Council and Oak Ridge Associated Universities. ROS acknowledges support from NSF grants PHYS-0121416 and PHYS-0353111.

#### REFERENCES

- Baker, J., Campanelli, M., Lousto, C. O., & Takahashi, R. 2002, *Phys. Rev. D*, 65, 124012
- Baker, J. G., Centrella, J., Choi, D., Koppitz, M., & Meter, J. v. 2006, submitted to PRD (gr-qc/0602026)
- Baumgardt, H., Hut, P., Makino, J., McMillan, S., & Portegies Zwart, S. 2003, *ApJ*, 582, L21
- Bender, P. L. & Hils, D. 1997, *Classical and Quantum Gravity*, 14, 1439
- Binney, J. & Tremaine, S. 1987, *Galactic dynamics* (Princeton, NJ, Princeton University Press, 1987, 747 p.)
- Blain, A. W., Kneib, J.-P., Ivison, R. J., & Smail, I. 1999, *ApJ*, 512, L87
- Campanelli, M., Lousto, C., & Zlochower, Y. 2006, *Phys. Rev. D*, 73, 061501
- Ebisuzaki, T., Makino, J., Tsuru, T. G., Funato, Y., Portegies Zwart, S., Hut, P., McMillan, S., Matsushita, S., Matsumoto, H., & Kawabe, R. 2001, *ApJ*, 562, L19
- Farmer, A. J. & Phinney, E. S. 2003, *MNRAS*, 346, 1197
- Flanagan, É. É. & Hughes, S. A. 1998, *Phys. Rev. D*, 57, 4535
- Freitag, M., Gürkan, M. A., & Rasio, F. A. 2006, *MNRAS*, 368, 141
- Gebhardt, K., Rich, R. M., & Ho, L. C. 2005, *ApJ*, 634, 1093
- Gerssen, J., van der Marel, R. P., Gebhardt, K., Guhathakurta, P., Peterson, R. C., & Pryor, C. 2002, *AJ*, 124, 3270
- Gürkan, M. A., Fregeau, J. M., & Rasio, F. A. 2006, *ApJ*, 640, L39
- Gürkan, M. A., Freitag, M., & Rasio, F. A. 2004, *ApJ*, 604, 632
- Hogg, D. W. 1999, (astro-ph/9905116)
- Hut, P., McMillan, S., Goodman, J., Mateo, M., Phinney, E. S., Pryor, C., Richer, H. B., Verbunt, F., & Weinberg, M. 1992, *PASP*, 104, 981
- Ivanova, N., Belczynski, K., Fregeau, J. M., & Rasio, F. A. 2005, *MNRAS*, 358, 572
- Joshi, K. J., Nave, C. P., & Rasio, F. A. 2001, *ApJ*, 550, 691
- Larson, S. L. 2003, Online Sensitivity Curve Generator, located at <http://www.srl.caltech.edu/~shane/sensitivity/>, based on Larson, S. L., Hiscock, W. A. & Hellings, R. W., 2000, *Phys. Rev. D*, 62, 062001
- Madau, P. & Pozzetti, L. 2000, *MNRAS*, 312, L9
- Miller, M. C. 2002, *ApJ*, 581, 438
- 2005, *ApJ*, 618, 426
- Miller, M. C. & Colbert, E. J. M. 2004, *International Journal of Modern Physics D*, 13, 1
- Porciani, C. & Madau, P. 2001, *ApJ*, 548, 522
- Portegies Zwart, S. F., Baumgardt, H., Hut, P., Makino, J., & McMillan, S. L. W. 2004, *Nature*, 428, 724
- Portegies Zwart, S. F., Makino, J., McMillan, S. L. W., & Hut, P. 1999, *A&A*, 348, 117
- Portegies Zwart, S. F. & McMillan, S. L. W. 2000, *ApJ*, 528, L17
- 2002, *ApJ*, 576, 899
- Quinlan, G. D. 1996, *New Astronomy*, 1, 35
- Spergel, D. N., Bean, R., Dore, O., Nolta, M. R., Bennett, C. L., Hinshaw, G., Jarosik, N., Komatsu, E., Page, L., Peiris, H. V., Verde, L., Barnes, C., Halpern, M., Hill, R. S., Kogut, A., Limon, M., Meyer, S. S., Odegard, N., Tucker, G. S., Weiland, J. L., Wollack, E., & Wright, E. L. 2006, submitted to *ApJ* (astro-ph/0603449)
- Steidel, C. C., Adelberger, K. L., Giavalisco, M., Dickinson, M., & Pettini, M. 1999, *ApJ*, 519, 1
- van der Marel, R. P., Gerssen, J., Guhathakurta, P., Peterson, R. C., & Gebhardt, K. 2002, *AJ*, 124, 3255
- Will, C. M. 2004, *ApJ*, 611, 1080

Yu, Q. & Tremaine, S. 2003, *ApJ*, 599, 1129  
Zhang, Q. & Fall, S. M. 1999, *ApJ*, 527, L81